

EXISTENCE RESULTS AND NUMERICAL SOLUTION OF FULLY FOURTH-ORDER NONLINEAR FUNCTIONAL DIFFERENTIAL EQUATIONS

DANG QUANG A¹, NGUYEN THANH HUONG^{2,*}

¹Center for Informatics and Computing, Vietnam Academy of Science and Technology,
18 Hoang Quoc Viet Street, Cau Giay District, Ha Noi, Viet Nam

²Thai Nguyen University of Sciences, Tan Thinh Ward, Thai Nguyen City,
Thai Nguyen Province, Viet Nam



Abstract. In this paper, we consider a boundary value problem for a fully fourth-order nonlinear functional differential equation which contains all lower derivatives of proportional delay arguments. By the reduction of the problem to operator equation for the right-hand side nonlinear function, we establish the existence and uniqueness of the solution and construct iterative methods on both continuous and discrete levels for solving it. We obtain the total error estimate for the discrete iterative solution. Many examples demonstrate the validity of the obtained theoretical results and the efficiency of the numerical method.

Keywords. Fully fourth-order nonlinear boundary value problem, functional differential equation, existence and uniqueness of solution, iterative method.

1. INTRODUCTION

In this paper, we consider a class of boundary value problems for fourth-order nonlinear functional differential equation

$$\begin{aligned} u''''(t) &= f(t, U(t)), \quad 0 < t < 1, \\ u(0) &= a, \quad u(1) = b, \quad u'(0) = c, \quad u'(1) = d, \end{aligned} \quad (1)$$

where $U(t) = (u(t), u(\varphi_0(t)), u'(t), u'(\varphi_1(t)), u''(t), u''(\varphi_2(t)), u'''(t), u'''(\varphi_3(t)))$, (2)

and $f : [0, 1] \times \mathbb{R}^8 \rightarrow \mathbb{R}$ and $\varphi_i : [0, 1] \rightarrow [0, 1]$ ($i = \overline{0, 3}$) are continuous functions.

A particular case of the problem (1) when $U(t) = (u(t), u(\varphi(t)))$ was studied in [1]. In this work, by reducing the problem to the equivalent integral equation, the authors established the existence and uniqueness of a solution under very strong assumptions, one of them is the Lipschitz condition of the function $f(t, u, v)$ in the variables u, v in the domain $[a, b] \times \mathbb{R}^2$. Under some other assumptions, the authors constructed an iterative scheme with the use of cubic spline interpolation at each iteration. The analysis of convergence was made, but it is a regret that in the proof of Theorem 8 [1, page 139], there are vital errors shown by us

*Corresponding author.

E-mail addresses: dangquanga@cic.vast.vn (D.Q. A); nguyenthanhhuong2806@gmail.com (N.T. Huong).

in a private letter to the authors and in [2]. After that, these errors were overcome in the corrigendum [3]. It should be emphasized that in all illustrative examples (Example 11 and Example 12 [1]), the conditions for ensuring the existence and uniqueness of solutions are not satisfied although the numerical results show good convergence.

In a very recent work [2], we proposed an effective method for investigating simultaneously the existence of solution and numerical method for the third-order functional differential equation

$$u'''(t) = f(t, u(t), u(\varphi(t))) \quad (3)$$

associated with general linear boundary conditions. The method is a further development of our method for nonlinear boundary value problems in [4, 5].

In connection to the general functional differential equation $u^{(p)}(t) = f(t, u(\varphi(t)))$, $p \geq 1$ it should be said that in its particular case when $\varphi(t) = \alpha t$, $0 < \alpha < 1$, the equation is called pantograph equation or proportional delay differential equation. This type of equation is intensively investigated in many works such as [6–10]. The above-mentioned works are concerned with initial value problems.

The solution of boundary value problems for second-order functional differential equation is considered in several works, e.g., [11–14], where numerical, semi-analytical, and neural network methods are used.

The present work aims to investigate the existence and uniqueness of the solution of the boundary value problem for general fourth-order nonlinear functional differential equation (1)-(2) and develop an efficient numerical method for finding the solution.

2. EXISTENCE AND UNIQUENESS OF SOLUTION

Following the approach in [2, 15] (see also [4, 5]) to investigate the problem (1)-(2), we introduce the nonlinear operator A defined in the space of continuous functions $C[0, 1]$ by the formula

$$(A\psi)(t) = f(t, U(t)), \quad (4)$$

where $u(t)$ is the solution of the problem

$$\begin{aligned} u''''(t) &= \psi(t), \quad 0 < t < 1, \\ u(0) &= a, \quad u(1) = b, \quad u'(0) = c, \quad u'(1) = d. \end{aligned} \quad (5)$$

It is easy to verify the following result.

Proposition 1. *If the function ψ is a fixed point of the operator A , i.e., ψ is the solution of the operator equation*

$$A\psi = \psi, \quad (6)$$

where A is defined by (4)-(5) then the function $u(t)$ determined from the BVP (5) is a solution of the BVP (1)-(2). Conversely, if the function $u(x)$ is the solution of the BVP (1)-(2) then the function

$$\psi(t) = f(t, U(t))$$

satisfies the operator equation (6).

Now, let $G(t, s)$ be the Green function of the problem (5). We have

$$G(t, s) = \frac{1}{6} \begin{cases} s^2(1-t)^2(3t-s-2ts), & 0 \leq s \leq t \leq 1, \\ t^2(1-s)^2(3s-t-2ts), & 0 \leq t \leq s \leq 1. \end{cases}$$

Denote $G_0(t, s) = G(t, s)$ and

$$G_i(t, s) = \frac{\partial^i G(t, s)}{\partial t^i}, \quad (i = \overline{1, 3}). \tag{7}$$

Then there hold the estimates

$$\int_0^1 |G_i(t, s)| ds \leq M_i, \quad (i = \overline{0, 3}) \tag{8}$$

with

$$M_0 = \frac{1}{384}, \quad M_1 = \frac{1}{72\sqrt{3}}, \quad M_2 = \frac{1}{12}, \quad M_3 = \frac{1}{2}. \tag{9}$$

The solution of the problem (5) is represented in the form

$$u(t) = g(t) + \int_0^1 G(t, s)\psi(s)ds, \tag{10}$$

where $g(t)$ is the polynomial of second degree satisfying the boundary conditions

$$g(0) = a, \quad g(1) = b, \quad g'(0) = c, \quad g'(1) = d. \tag{11}$$

Taking the derivatives of (9), we obtain

$$u^{(i)}(t) = g^{(i)}(t) + \int_0^1 G_i(t, s)\psi(s)ds, \quad (i = \overline{0, 3}). \tag{12}$$

Therefore, from (7), we have

$$\|u^{(i)}\| \leq \|g^{(i)}\| + M_i\|\psi\|, \quad (i = \overline{0, 3}), \tag{13}$$

where $\|\cdot\|$ is the norm in the space $C[0, 1]$, $\|v\| = \max_{0 \leq t \leq 1} |v(t)|$, $v \in C[0, 1]$.

Now for every positive number M define the domain

$$\mathcal{D}_M = \left\{ (t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z}) \mid 0 \leq t \leq 1; |u|, |\bar{u}| \leq \|g\| + M_0M; |y|, |\bar{y}| \leq \|g'\| + M_1M, \right. \\ \left. |v|, |\bar{v}| \leq \|g''\| + M_2M; |z|, |\bar{z}| \leq \|g'''\| + M_3M \right\}. \tag{14}$$

As usual, the closed ball of the radius M centered at 0 in the space of continuous functions $C[0, 1]$ is denoted by $B[0, M]$.

Theorem 1. *Assume that:*

- (i) *The function $\varphi_k(t)$ ($k = \overline{0, 3}$) are continuous functions from $[0, 1]$ to $[0, 1]$.*
- (ii) *The function $f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})$ is continuous and bounded by M in the domain \mathcal{D}_M , i.e.,*

$$|f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})| \leq M, \quad \forall (t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z}) \in \mathcal{D}_M. \tag{15}$$

- (iii) *The function $f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})$ satisfies the Lipschitz conditions in the variables $u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z}$ with the coefficients $L_i \geq 0$, ($i = \overline{0, 7}$) in \mathcal{D}_M , i.e.,*

$$|f(t, u_2, \bar{u}_2, y_2, \bar{y}_2, v_2, \bar{v}_2, z_2, \bar{z}_2) - f(t, u_1, \bar{u}_1, y_1, \bar{y}_1, v_1, \bar{v}_1, z_1, \bar{z}_1)| \\ \leq L_0|u_2 - u_1| + L_1|\bar{u}_2 - \bar{u}_1| + L_2|y_2 - y_1| + L_3|\bar{y}_2 - \bar{y}_1| \\ + L_4|v_2 - v_1| + L_5|\bar{v}_2 - \bar{v}_1| + L_6|z_2 - z_1| + L_7|\bar{z}_2 - \bar{z}_1|, \\ \forall (t, u_i, \bar{u}_i, y_i, \bar{y}_i, v_i, \bar{v}_i, z_i, \bar{z}_i) \in \mathcal{D}_M \quad (i = \overline{1, 2}). \tag{16}$$

$$(iv) \quad q := (L_0 + L_1)M_0 + (L_2 + L_3)M_1 + (L_4 + L_5)M_2 + (L_6 + L_7)M_3 < 1. \quad (17)$$

The the problem (1)-(2) has a unique solution $u(t) \in C^4[0, 1]$, satisfying the estimate

$$|u^{(i)}(t)| \leq \|g^{(i)}\| + M_i M, (i = \overline{0, 3}) \quad \forall t \in [0, 1]. \quad (18)$$

Proof. Sketch the steps of the proof:

i) Show that the operator A is a mapping $B[0, M] \rightarrow B[0, M]$.

ii) Show that A is a contraction mapping in $B[0, M]$.

Then the operator equation (6) has a unique solution $\psi \in B[0, M]$. By Proposition 1 the solution of the problem (5) for this right-hand side $\psi(t)$ is the solution of the original problem (1)-(2).

The detailed proof is similar to that in [2] for the equation (3). ■

3. SOLUTION METHOD AND ITS CONVERGENCE

Consider the following iterative method:

1. Given $\psi_0 \in B[0, M]$, for example,

$$\psi_0(t) = f(t, 0, 0, 0, 0, 0, 0, 0). \quad (19)$$

2. Knowing $\psi_k(t)$ ($k = 0, 1, \dots$) compute

$$\begin{aligned} u_k(t) &= g(t) + \int_0^1 G(t, s)\psi_k(s)ds, \\ \bar{u}_k(t) &= g(\varphi_0(t)) + \int_0^1 G(\varphi_0(t), s)\psi_k(s)ds, \\ y_k(t) &= g'(t) + \int_0^1 G_1(t, s)\psi_k(s)ds, \\ \bar{y}_k(t) &= g'(\varphi_1(t)) + \int_0^1 G_1(\varphi_1(t), s)\psi_k(s)ds, \\ v_k(t) &= g''(t) + \int_0^1 G_2(t, s)\psi_k(s)ds, \\ \bar{v}_k(t) &= g''(\varphi_2(t)) + \int_0^1 G_2(\varphi_2(t), s)\psi_k(s)ds, \\ z_k(t) &= g'''(t) + \int_0^1 G_3(t, s)\psi_k(s)ds, \\ \bar{z}_k(t) &= g'''(\varphi_3(t)) + \int_0^1 G_3(\varphi_3(t), s)\psi_k(s)ds. \end{aligned} \quad (20)$$

3. Update

$$\psi_{k+1}(t) = f(t, u_k(t), \bar{u}_k(t), y_k(t), \bar{y}_k(t), v_k(t), \bar{v}_k(t), z_k(t), \bar{z}_k(t)). \quad (21)$$

Set

$$p_k = \frac{q^k}{1 - q}, \quad d = \|\psi_1 - \psi_0\|. \quad (22)$$

Theorem 2. [Convergence] *Under the assumptions of Theorem 1 the above iterative method converges and there holds the estimate*

$$\|u_k^{(i)} - u^{(i)}\| \leq M_i p_k d, \quad (i = \overline{0, 3}),$$

where u is the exact solution of the problem (1)-(2), $u^{(i)}$ is its derivative of order i and M_i is given by (9).

This theorem follows straightforward by from the convergence of the iterative method for a fixed point of the operator A , the representations (10), (12), and the formulas for computing $u_k(t)$, $y_k(t)$, $v_k(t)$, $z_k(t)$ in (20).

Now we design a discrete iterative scheme for realizing the above iterative method on a continuous level. For this purpose, we construct the uniform grid $\bar{\omega}_h = \{t_i = ih, h = a/N, i = 0, 1, \dots, N\}$ on the interval $[0, 1]$ and denote by $\Phi_k(t)$, $U_k(t)$, $\bar{U}_k(t)$, $Y_k(t)$, $\bar{Y}_k(t)$, $V_k(t)$, $\bar{V}_k(t)$, $Z_k(t)$, $\bar{Z}_k(t)$ the grid functions, which are defined on the grid $\bar{\omega}_h$ and approximate the functions $\psi_k(t)$, $u_k(t)$, $\bar{u}_k(t)$, $y_k(t)$, $\bar{y}_k(t)$, $v_k(t)$, $\bar{v}_k(t)$, $z_k(t)$, $\bar{z}_k(t)$ on this grid, respectively.

Below are the steps of the discrete iterative method:

1. Given
$$\Psi_0(t_i) = f(t_i, 0, 0, 0, 0, 0, 0, 0, 0), \quad i = 0, \dots, N. \tag{23}$$

2. Knowing $\Psi_k(t_i)$, $k = 0, 1, \dots$; $i = 0, \dots, N$, compute approximately the definite integrals (20) by the trapezoidal rule

$$\begin{aligned} U_k(t_i) &= g(t_i) + \sum_{j=0}^N h\rho_j G(t_i, t_j) \Psi_k(t_j), \\ \bar{U}_k(t_i) &= g(\xi_{0i}) + \sum_{j=0}^N h\rho_j G(\xi_{0i}, t_j) \Psi_k(t_j), \\ Y_k(t_i) &= g'(t_i) + \sum_{j=0}^N h\rho_j G_1(t_i, t_j) \Psi_k(t_j), \\ \bar{Y}_k(t_i) &= g'(\xi_{1i}) + \sum_{j=0}^N h\rho_j G_1(\xi_{1i}, t_j) \Psi_k(t_j), \\ V_k(t_i) &= g''(t_i) + \sum_{j=0}^N h\rho_j G_2(t_i, t_j) \Psi_k(t_j), \\ \bar{V}_k(t_i) &= g''(\xi_{2i}) + \sum_{j=0}^N h\rho_j G_2(\xi_{2i}, t_j) \Psi_k(t_j), \\ Z_k(t_i) &= g'''(t_i) + \sum_{j=0}^N h\rho_j G_3^*(t_i, t_j) \Psi_k(t_j), \\ \bar{Z}_k(t_i) &= g'''(\xi_{3i}) + \sum_{j=0}^N h\rho_j G_3^*(\xi_{3i}, t_j) \Psi_k(t_j), \quad (i = 0, \dots, N), \end{aligned} \tag{24}$$

where ρ_j are the weights

$$\rho_j = \begin{cases} 1/2, & j = 0, N \\ 1, & j = 1, 2, \dots, N - 1, \end{cases} \quad \xi_{mi} = \varphi_m(t_i), \quad m = \overline{0, 3}$$

and

$$G_3^*(t, s) = \begin{cases} G_3(t, s), & s \neq t, \\ \frac{1}{2}[\lim_{s \rightarrow t-0} G_3(t, s) + \lim_{s \rightarrow t+0} G_3(t, s)], & s = t. \end{cases}$$

3. Update

$$\Psi_{k+1}(t_i) = f(t_i, U_k(t_i), \bar{U}_k(t_i), Y_k(t_i), \bar{Y}_k(t_i), V_k(t_i), \bar{V}_k(t_i), Z_k(t_i), \bar{Z}_k(t_i)). \tag{25}$$

To study the convergence of the above discrete iterative method, we need some auxiliary results.

Proposition 2. *Assume that the functions $\varphi_m(t)$, ($m = \overline{0, 3}$) have continuous derivatives up to second order in $[0, 1]$ and the function $f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})$ has all partial derivatives continuous up to second order in the domain \mathcal{D}_M . Then the functions generated by the iterative method (19)-(21) satisfy the following conditions*

$$\begin{aligned} z_k(t) \in C^3[0, 1], \quad v_k(t) \in C^4[0, 1], \quad y_k(t) \in C^5[0, 1], \quad u_k(t) \in C^6[0, 1], \\ \psi_k(t), \quad \bar{u}_k(t), \quad \bar{y}_k(t), \quad \bar{v}_k(t), \quad \bar{z}_k(t) \in C^2[0, 1]. \end{aligned} \tag{26}$$

Proof. We shall prove the above proposition by induction for k .

For $k = 0$, by the assumption on the function f , we have $\psi_0(t) = f(t, 0, 0, 0, 0, 0, 0, 0, 0) \in C^2[0, 1]$. Taking into account the expression of $G_3(t, s)$ (see Appendix), we have

$$\begin{aligned} z_0(t) &= g'''(t) + \int_0^1 G_3(t, s)\psi_0(s)ds \\ &= g'''(t) + \int_0^1 s^2(3 - 2s)\psi_0(s)ds - \int_t^1 \psi_0(s)ds. \end{aligned}$$

Therefore, $z'_0(t) = \psi_0(t)$. It implies that $z'_0(t) \in C^2[0, 1]$ and consequently, $z_0(t) \in C^3[0, 1]$. In view of (20) it follows that $v_0(t) \in C^4[0, 1]$, $y_0(t) \in C^5[0, 1]$, $u_0(t) \in C^6[0, 1]$. Next, we can obtain $\bar{z}'_0(t) = \varphi'_3(t)\psi_0(\varphi_3(t)) \in C^1[0, 1]$. Hence, $\bar{z}_0(t) \in C^2[0, 1]$. It is also easy to show that $\bar{v}_0(t), \bar{y}_0(t), \bar{u}_0(t) \in C^2[0, 1]$.

Thus, the proposition is true for $k = 0$.

Now suppose that the proposition is true for $k \geq 0$, i.e., there holds (26). Then, from the assumption of the smoothness of the function f and from the formula (21), it follows that $\psi_{k+1}(t) \in C^2[0, 1]$. Making a similar argument as for the case $k = 0$, we obtain

$$\begin{aligned} z_{k+1}(t) \in C^3[0, 1], \quad v_{k+1}(t) \in C^4[0, 1], \quad y_{k+1}(t) \in C^5[0, 1], \quad u_{k+1}(t) \in C^6[0, 1], \\ \bar{u}_{k+1}(t), \bar{y}_{k+1}(t), \bar{v}_{k+1}(t), \bar{z}_{k+1}(t) \in C^2[0, 1]. \end{aligned}$$

Thus, the proposition is proved. ■

Proposition 3. For any function $\psi(t) \in C^2[0, 1]$ there hold the estimates

$$\int_0^1 G_m(t_i, s)\psi(s)ds = \sum_{j=0}^N h\rho_j G_m(t_i, s_j)\psi(s_j) + O(h^2), \quad m = 0, 1, 2, \quad (27)$$

$$\int_0^1 G_3(t_i, s)\psi(s)ds = \sum_{j=0}^N h\rho_j G_3^*(t_i, s_j)\psi(s_j) + O(h^2), \quad (28)$$

$$\int_0^1 G_m(\xi_{mi}, s)\psi(s)ds = \sum_{j=0}^N h\rho_j G_m(\xi_{mi}, s_j)\psi(s_j) + O(h^2), \quad m = 0, 1, 2, \quad (29)$$

$$\int_0^1 G_3(\xi_{3i}, s)\psi(s)ds = \sum_{j=0}^N h\rho_j G_3^*(\xi_{3i}, s_j)\psi(s_j) + O(h^2), \quad (30)$$

where $\xi_{mi} = \varphi_m(t_i)$, $m = \overline{0, 3}$.

Proof. For proof of (27)-(28) see [16, Proposition 2], and for (29)-(30) see [2, Proposition 3.3]. ■

Proposition 4. Under the assumptions of Theorem 1, we have the estimates

$$\|\Psi_k - \psi_k\|_{\bar{\omega}_h} = O(h^2), \quad \|U_k - u_k\|_{\bar{\omega}_h} = O(h^2), \quad (31)$$

$$\|Y_k - y_k\|_{\bar{\omega}_h} = O(h^2), \quad \|V_k - v_k\|_{\bar{\omega}_h} = O(h^2), \quad \|Z_k - z_k\|_{\bar{\omega}_h} = O(h^2), \quad (32)$$

where $\|\cdot\|_{\bar{\omega}_h}$ is the max-norm of the grid function defined on the grid $\bar{\omega}_h$.

Proof. It is not hard to prove the proposition by induction using Propositions 2 and 3, and taking into account the formulas (19), (23) and (20), (24) of the iterative methods on continuous and on discrete levels. ■

Similar but simpler propositions are proved in detail in [2, Proposition 3.5] and [16, Proposition 3].

Now combining Proposition 4 with Theorem 2, we obtain the following result on the error estimate of the actually obtained numerical solution of the original problem.

Theorem 3. Under the assumptions of Theorem 1 for the approximate solution of the problem (1)-(2) obtained by the discrete iterative method (23)-(25), we have the estimates

$$\begin{aligned} \|U_k - u\|_{\bar{\omega}_h} &\leq M_0 p_k d + O(h^2), \quad \|Y_k - u'\|_{\bar{\omega}_h} \leq M_1 p_k d + O(h^2), \\ \|V_k - u''\|_{\bar{\omega}_h} &\leq M_2 p_k d + O(h^2), \quad \|Z_k - u'''\|_{\bar{\omega}_h} \leq M_3 p_k d + O(h^2), \end{aligned}$$

where p_k and d are defined by (22).

4. EXAMPLES

In all examples below first, we verify the conditions of Theorem 1 which guarantees the existence and uniqueness of a solution and also the convergence of the iterative methods. After that, we perform the iterative method, (23)-(25) until $\|U_k - U_{k-1}\|_{\bar{\omega}_h} \leq 10^{-16}$. In the

tables of results for the convergence of the iterative method $N + 1$ is the number of grid points, $h = 1/N$, K is the number of performed iterations, u is the exact solution if it is known and $Error = \|U_K - u\|_{\bar{\omega}_h}$ and $Error1 = \|Y_K - u'\|_{\bar{\omega}_h}$ are errors of the solution and its first derivative, respectively.

Example 1. [1, Example 11]. Consider the following problem

$$\begin{aligned} u^{(4)}(t) &= \frac{22}{(t+1)^5} + \frac{1}{(t+1)^2} \left([u(t)]^2 + [u(t)]^3 \right) u\left(\frac{t}{2}\right), \quad 0 < t < 1, \\ u(0) &= 1, \quad u(1) = \frac{1}{2}, \quad u'(0) = -1, \quad u'(1) = -\frac{1}{4}, \end{aligned} \quad (33)$$

for which the exact solution is $u(t) = \frac{1}{t+1}$.

For this problem, the right-hand side function f is

$$f(t, u, \bar{u}) = \frac{22}{(t+1)^5} + \frac{1}{(t+1)^2} (u^2 + u^3) \bar{u}.$$

Obviously, this function does not satisfy the Lipschitz conditions in the domain $[0, 1] \times \mathbb{R}^2$. Thus, the important condition (ii) in [1, page 131] is not met. So, the uniqueness of the solution is not ensured, and of course, Theorem 2 and Theorem 8 there on the convergence of the iterative method are not applicable. Differently from Bica [1] below, we show that the problem (33) has a unique solution and the discrete iterative method (23)-(25) converges with accuracy of second order. Indeed, for the problem it is easy to find the function $g(t) = -\frac{1}{4}t^3 + \frac{3}{4}t^2 - t + 1$ satisfying the boundary conditions, and $\|g\| = \|g'\| = 1$, $\|g''\| = \|g'''\| = \frac{3}{2}$. It is possible to verify that for $M = 25$, we have $|f(t, u, \bar{u})| \leq M$ in the domain $\mathcal{D}_M = \left\{ (t, u, \bar{u}) \mid 0 \leq t \leq 1; |u|, |\bar{u}| \leq 1 + M_0 M \right\}$. Further, in \mathcal{D}_M the function $f(t, u, \bar{u})$ satisfies the Lipschitz conditions in u and \bar{u} with the coefficients $L_0 = 6$, $L_1 = 2.4$. Therefore, $q = (L_0 + L_1)M_0 = 0.0219 < 1$. By Theorem 1 the problem (33) has a unique solution and by Theorem 3 the iterative method (23)-(25) converges.

In Table 1 below, the convergence of the iterative method is reported. As shown in the

Table 1: The convergence in Example 1

N	h^2	K	$Error$	$Error1$
50	4.0000e-04	8	1.4520e-08	3.8031e-08
100	1.0000e-04	8	9.0870e-10	2.3801e-09
150	4.4444e-05	8	1.7954e-10	4.7028e-10
200	2.5000e-05	8	5.6812e-11	1.4881e-10
300	1.1111e-05	9	1.1223e-11	2.9396e-11
400	6.2500e-06	9	3.5512e-12	9.3012e-12
500	4.0000e-06	8	1.4546e-12	3.8097e-12
800	1.5625e-06	9	2.2204e-13	5.8115e-13
1000	1.0000e-06	9	9.1038e-14	2.3798e-13

table, the errors of the approximate solution and its first derivative are of order 4 although Theorem 3 ensures only order 2. This fact also will be observed in the next example.

Example 2. [1, Example 12]. Consider the following problem

$$\begin{aligned}
 u^{(4)}(t) &= e^{-t}[u(t)]^{3/2}u\left(\frac{t}{2}\right), \quad t \in (0, 1), \\
 u(0) &= 1, \quad u(1) = e, \quad u'(0) = 1, \quad u'(1) = e,
 \end{aligned}
 \tag{34}$$

for which the exact solution is $u(t) = e^t$.

For this problem the right-hand side function f is

$$f(t, u, \bar{u}) = e^{-t}u^{3/2}\bar{u}.$$

As in the previous example, the function $f(t, u, \bar{u})$ does not satisfy the Lipschitz conditions in the domain $[0, 1] \times \mathbb{R}^2$. So, the important condition (ii) in [1, page 131] is not met. Hence, the uniqueness of the solution is not ensured, and of course, Theorem 2 and Theorem 8 there on the convergence of the iterative method are not applicable. Below, the theory in the previous section is applied to the example. We have for this example $g(t) = (3 - e)t^3 + (2e - 5)t^2 + t + 1$, and consequently, $\|g\| = e$. It is possible to verify that for $M = 15$, we have $|f(t, u, \bar{u})| \leq M$ in the domain

$$\mathcal{D}_M = \left\{ (t, u, \bar{u}) \mid 0 \leq t \leq 1; |u|, |\bar{u}| \leq e + M_0M \right\}.$$

Further, in \mathcal{D}_M the function $f(t, u, \bar{u})$ satisfies the Lipschitz conditions in u and \bar{u} with the coefficients $L_0 = 7, L_1 = 5$. Therefore, $q = (L_0 + L_1)M_0 = 0.0313 < 1$. By Theorem 1, the problem (33) has a unique solution and by Theorem 3, the iterative method (23)-(25) converges.

The results of the convergence of the iterative method for this example are given in Table 2.

Table 2: The convergence in Example 2

N	h^2	K	$Error$	$Error1$
50	4.0000e-04	9	3.0102e-10	2.2923e-09
100	1.0000e-04	9	1.8814e-11	1.4328e-10
150	4.4444e-05	9	3.7157e-12	2.8302e-11
200	2.5000e-05	10	1.1755e-12	8.9548e-12
300	1.1111e-05	10	2.3226e-13	1.7686e-12
400	6.2500e-06	10	7.3275e-14	5.5933e-13
500	4.0000e-06	9	3.0198e-14	2.2893e-13
800	1.5625e-06	9	4.8850e-15	3.4639e-14
1000	1.0000e-06	9	2.2204e-15	1.4211e-14

Example 3. Consider a more complicated problem

$$\begin{aligned}
 u^{(4)}(t) &= e^t + \frac{1}{9} \left([u\left(\frac{t}{2}\right)]^2 u'''\left(\frac{t}{2}\right) - u'(t)u''\left(\frac{t}{2}\right) + u(t)u'''(t) - [u''(t)]^2 \right), \quad t \in (0, 1), \\
 u(0) &= 1, \quad u(1) = e, \quad u'(0) = 1, \quad u'(1) = e,
 \end{aligned}
 \tag{35}$$

for which the exact solution is $u(t) = e^t$.

For this example

$$f = f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z}) = e^t + \frac{1}{9}(\bar{u}^2\bar{z} - y\bar{v} + uz - v^2),$$

$$g(t) = (3 - e)t^3 + (2e - 5)t^2 + t + 1.$$

So, we have $\|g\| = \|g'\| = e$, $\|g''\| = 8 - 2e$, $\|g'''\| = 18 - 6e$. It is possible to check that for $M = 20$ there holds $|f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})| \leq M$ in \mathcal{D}_M , where \mathcal{D}_M is defined by (14). In this domain, the function $f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})$ satisfies the Lipschitz conditions in the variables $u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z}$ with the coefficients $L_0 = 1.30$, $L_1 = 7.20$, $L_2 = 0.47$, $L_3 = 0$, $L_4 = 0.94$, $L_5 = 0.32$, $L_6 = 0.31$, and $L_7 = 0.86$. Therefore, $q = 0.6446 < 1$ and the problem (35) has a unique solution, and the iterative method (23)-(25) converges. The results of the convergence of the problem are given in Table 3.

Table 3: The convergence in Example 3

N	h^2	K	$Error$	$Error1$
50	4.0000e-04	9	9.2553e-08	3.0639e-07
100	1.0000e-04	9	2.3182e-08	7.6420e-08
150	4.4444e-05	9	1.0307e-08	3.3946e-08
200	2.5000e-05	11	5.7979e-09	1.9088e-08
300	1.1111e-05	9	2.5771e-09	8.4831e-09
400	6.2500e-06	9	1.4497e-09	4.7715e-09
500	4.0000e-06	9	9.2781e-10	3.0537e-09
800	1.5625e-06	8	3.6243e-10	1.1928e-09
1000	1.0000e-06	8	2.3196e-10	7.6339e-10

It is easy to see in Table 3 that for this example when the right-hand side is rather complicated, the errors of the approximate solution and its first derivative are not of order 4 as in the previous examples (Bica's examples), where the right-hand sides are simple. Obviously, here $Error$ and $Error1$ are of $O(h^3)$.

Example 4. Consider the following problem

$$u^{(4)}(t) = t^2 - \frac{1}{4}u(t) + \frac{1}{4}[u(\varphi_0(t))]^2 + \frac{1}{2}u'(t)u'(\varphi_1(t)) + \frac{1}{8}[u''(t) + u''(\varphi_2(t))]u(t)$$

$$+ \frac{1}{4}[\sin(u'''(t)) + \cos(u'''(\varphi_3(t)))], \quad t \in (0, 1), \quad (36)$$

$$u(0) = 1, \quad u(1) = \frac{19}{6}, \quad u'(0) = 1, \quad u'(1) = \frac{7}{2},$$

where $\varphi_0(t) = \frac{t}{2}$, $\varphi_1(t) = t^2$, $\varphi_2 = \frac{t^2}{2}$, $\varphi_3(t) = \frac{t^2}{3}$. At this moment, we do not know any information on the solvability of the problem.

For the above problem

$$f = t^2 - \frac{1}{4}u + \frac{1}{4}\bar{u}^2 + \frac{1}{2}y\bar{y} + \frac{1}{8}(v + \bar{v})u + \frac{1}{4}(\sin(z) + \cos(\bar{z})),$$

$$g(t) = \frac{t^3}{6} + t^2 + t + 1.$$

Hence, $\|g\| = \frac{19}{6}$, $\|g'\| = \frac{7}{2}$, $\|g''\| = 3$, $\|g'''\| = 1$. It is possible to verify that with $M = 23$ there holds $|f| \leq M$ in the domain \mathcal{D}_M , where \mathcal{D}_M is defined by (14). Also, in \mathcal{D}_M the function $f(t, u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z})$ satisfies the Lipschitz conditions in the variables $u, \bar{u}, y, \bar{y}, v, \bar{v}, z, \bar{z}$ with the coefficients $L_0 = 1.48$, $L_1 = 1.62$, $L_2 = L_3 = 3.7$, $L_4 = L_5 = 0.41$, $L_6 = L_7 = 0.25$. Therefore, $q = 0.3857 < 1$ and the problem (36) has a unique solution and the iterative method (23)-(25) converges. The results of the convergence of the problem are given in Table 4. Recall that in the table, K is the number of iterations performed when $\|U_k - U_{k-1}\|_{\bar{\omega}_h} \leq 10^{-16}$.

Table 4: The convergence in Example 4

N	K
50	12
100	13
150	13
200	12
300	13
400	14
500	14
800	13
1000	13

The graph of the found approximate solution is depicted in Figure 1.




Figure 1: The graph of the approximate solution in Example 4

Remark. It should be remarked that Theorem 1 and Theorem 3 give only sufficient conditions for the existence and uniqueness of the solution, and the convergence of the iterative method for finding the solution. In the case if these conditions are not satisfied, the iterative method also can be convergent. Below, we give some examples to illustrate this remark.

Example 5. Consider the following problem

$$\begin{aligned} u^{(4)}(t) &= [u''(\frac{t}{4})]^4, \quad 0 < t < 1, \\ u(0) &= 1, \quad u(1) = e, \quad u'(0) = 1, \quad u'(1) = e, \end{aligned} \quad (37)$$

which has the exact solution $u(t) = e^t$.

For this example, $f = f(t, \bar{v}) = (\bar{v})^4$. It is impossible to find $M > 0$ so that $|f| \leq M$ in the domain $\mathcal{D}_M = \{(t, \bar{v}) \mid 0 \leq t \leq 1, |\bar{v}| \leq 8 - 2e + \frac{M}{12}\}$. Recall that $8 - 2e = \|g''\|$ (see Example 3). Therefore, the theorems 1 and 2 are not applicable. Regardless of this, the iterative method (23)-(25) converges to the exact solution $u(t) = e^t$. The results of convergence for the problem (37) are shown in Table 5.

Table 5: The convergence in Example 5

N	h^2	K	$Error$	$Error1$
50	4.0000e-04	20	4.3691e-08	1.3415e-07
100	1.0000e-04	21	1.0868e-08	3.3749e-08
150	4.4444e-05	20	4.8264e-09	1.5026e-08
200	2.5000e-05	21	2.7140e-09	8.4562e-09
300	1.1111e-05	22	1.2059e-09	3.7597e-09
400	6.2500e-06	22	6.7828e-10	2.1150e-09
500	4.0000e-06	22	4.3409e-10	1.3537e-09
800	1.5625e-06	22	1.6956e-10	5.2882e-10
1000	1.0000e-06	22	1.0852e-10	3.3845e-10

Example 6. Consider the following problem

$$\begin{aligned} u^{(4)}(t) &= (u(t))^2 + (u''(\frac{t^2}{2}))^4, \quad t \in (0, 1), \\ u(0) &= 1, \quad u(1) = \frac{19}{6}, \quad u'(0) = 1, \quad u'(1) = \frac{7}{2}. \end{aligned} \quad (38)$$

We do not know any information about the solution to the problem. And it is easy to verify that the conditions of Theorem 1 are not satisfied. Therefore, it is not ensured of the existence of solutions and the convergence of the iterative method. However, the results of numerical experiments show that the iterative method (23)- (25) has a good convergence. The number of iterations needed for achieving the tolerance $\|U_k - U_{k-1}\|_{\bar{\omega}_h} \leq 10^{-16}$ is 15 or 16.

5. CONCLUSION

In this paper, we have considered the Dirichlet problem for a fully fourth-order nonlinear functional differential equation, where its right-hand side contains all lower derivatives of proportional delay arguments. By reducing the problem to an operator equation for the right-hand side nonlinear function, we have established the existence and uniqueness of the solution under some easily verified conditions and constructed an efficient iterative method

for finding the solution at both continuous and discrete levels. A total error of the actual numerical solution which consists of the error of the iterative process and the error of discretization at each iteration is obtained. Many examples with numerical experiments confirm the applicability and validity of the obtained theoretical results. To our best knowledge, this is the first time a boundary value problem for a fully fourth-order nonlinear functional differential equation is studied.

It should be said that the proposed approach here to boundary value problems for nonlinear functional differential equations is very effective because it simultaneously gives the existence results and the convergence of the solution method which is very easily realized. The approach can be applied to a wide class of boundary value problems for nonlinear functional differential equations of arbitrary order and any linear boundary conditions.

ACKNOWLEDGMENT

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 102.01-2021.03.

APPENDIX

Derivatives in t of the Green function $G(t, s)$:

$$G(t, s) = \frac{1}{6} \begin{cases} s^2(1-t)^2(3t-s-2ts), & 0 \leq s \leq t \leq 1, \\ t^2(1-s)^2(3s-t-2ts), & 0 \leq t \leq s \leq 1, \end{cases}$$

$$G_1(t, s) = \frac{\partial G(t, s)}{\partial t} = \begin{cases} -(s^2(2t-2)(s-3t+2st))/6 - (s^2(2s-3)(t-1)^2)/6, & 0 \leq s \leq t \leq 1, \\ -(t^2(2s+1)(s-1)^2)/6 - (t(s-1)^2(t-3s+2st))/3, & 0 \leq t \leq s \leq 1, \end{cases}$$

$$G_2(t, s) = \frac{\partial^2 G(t, s)}{\partial t^2} = \begin{cases} -(s^2(s-3t+2st))/3 - (s^2(2s-3)(2t-2))/3, & 0 \leq s \leq t \leq 1, \\ -((s-1)^2(t-3s+2st))/3 - (2t(2s+1)(s-1)^2)/3, & 0 \leq t \leq s \leq 1, \end{cases}$$

$$G_3(t, s) = \frac{\partial^3 G(t, s)}{\partial t^3} = \begin{cases} s^2(3-2s), & 0 \leq s < t \leq 1, \\ s^2(3-2s) - 1, & 0 \leq t < s \leq 1, \end{cases}$$

$$G_3^*(t, s) = \begin{cases} s^2(3-2s), & 0 \leq s < t \leq 1, \\ t^2(3-2t) - 1/2, & s = t, \\ s^2(3-2s) - 1, & 0 \leq t < s \leq 1. \end{cases}$$

REFERENCES

- [1] A. M. Bica, M. Curila, and S. Curila, "Two-point boundary value problems associated to functional differential equations of even order solved by iterated splines," *Applied Numerical Mathematics*, vol. 110, pp. 128–147, 2016.
- [2] Q. A. Dang and Q. L. Dang, "A unified approach to study the existence and numerical solution of functional differential equation," *Applied Numerical Mathematics*, vol. 170, pp. 208–218, 2021.
- [3] A. M. Bica, "Corrigendum to "two-point boundary value problems associated to functional differential equations of even order solved by iterated splines" [appl. numer. math. 110(2016) 128-147]," *Applied Numerical Mathematics*, vol. 165, pp. 620–621, 2021.

- [4] Q. A. Dang and T. K. Q. Ngo, “Existence results and iterative method for solving the cantilever beam equation with fully nonlinear term,” *Nonlinear Analysis: Real World Applications*, vol. 36, pp. 56–68, 2017.
- [5] Q. A. Dang, T. H. Nguyen, and H. H. Truong, “Unified approach to fully fourth order nonlinear problems,” *Azerbaijan Journal of Mathematics*, vol. 9, no. 2, pp. 114–132, 2019.
- [6] M. Chamekh, T. M. Elzaki, and N. Brik, “Semi-analytical solution for some proportional delay differential equations,” *SN Applied Sciences*, vol. 1:148, 2019. [Online]. Available: <https://doi.org/10.1007/s42452-018-0130-8>
- [7] H. Jafari, M. Mahmoudi, and M. H. N. Skandari, “A new numerical method to solve pantograph delay differential equations with convergence analysis,” *Advances in Continuous and Discrete Models*, no. 129, 2021. [Online]. Available: <https://doi.org/10.1186/s13662-021-03293-0>
- [8] M. A. Z. Raja, “Numerical treatment for boundary value problems of pantograph functional differential equation using computational intelligence algorithms,” *Applied Soft Computing*, vol. 24, pp. 806–821, 2014.
- [9] S. Y. Reutskiy, “A new collocation method for approximate solution of the pantograph functional differential equations with proportional delay,” *Applied Mathematics and Computation*, vol. 266, pp. 642–655, 2015.
- [10] S. Yüzbasi and M. Karacayir, “A galerkin-like approach to solve multi-pantograph type delay differential equations,” *Filomat*, vol. 32, pp. 409–422, 2018.
- [11] A. M. Bica, M. Curila, and S. Curila, “About a numerical method of successive interpolations for two-point boundary value problems with deviating argument,” *Applied Mathematics and Computation*, vol. 217, pp. 7772–7789, 2011.
- [12] C.-C. Hou, T. E. Simos, and I. T. Famelis, “Neural network solution of pantograph type differential equations,” *Mathematical Methods in the Applied Sciences*, pp. 1–6, 2020. [Online]. Available: <https://doi.org/10.1002/mma.6126>
- [13] S. A. Khuri and A. Sayfy, “Numerical solution of functional differential equations: a green’s function-based iterative approach,” *International Journal of Computer Mathematics*, vol. 95, pp. 1937–1949, 2018.
- [14] A. M. Wazwaz, M. A. Z. Raja, and M. I. Syam, “Reliable treatment for solving boundary value problems of pantograph delay differential equations,” *Romanian Reports in Physics*, vol. 69, no. 102, pp. 1–15, 2017.
- [15] Q. A. Dang and Q. L. Dang, “Existence results and numerical method for solving a fourth-order nonlinear integro-differential equation,” *Numerical Algorithms*, vol. 90, pp. 563–576, 2022.
- [16] —, “Existence results and iterative method for a fully fourth-order nonlinear integral boundary value problem,” *Numerical Algorithms*, vol. 85, pp. 887–907, 2020.

Received on September 06, 2023

Accepted on December 18, 2023