

**TREE SNAILS *Amphidromus* Albers, 1850 (Pulmonata: Camaenidae)
COLLECTED IN VIETNAM**

**Do Van Nhuong¹, Dinh Dieu Thuy^{2,3,*}, Do Duc Sang⁴, Nguyen Thanh Tung⁵,
Le Hung Anh^{3,6}, Arthur E. Bogan⁷, Do Van Tu^{3,6}**

¹Ha Noi National University of Education, Vietnam National University, Ha Noi, Vietnam

²Vietnam National Museum of Nature, Ha Noi, Vietnam

³Graduate University of Science and Technology, Vietnam Academy of Science and
Technology, 18 Hoang Quoc Viet, Ha Noi, Vietnam

⁴Ha Noi University of Science, VNU, 334 Nguyen Trai, Ha Noi, Vietnam

⁵Can Tho University, Can Tho, Vietnam

⁶Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology,
18 Hoang Quoc Viet, Ha Noi, Vietnam

⁷North Carolina Museum of Natural Sciences 11 West Jones St. Raleigh, NC 27601

Received 14 September 2023; accepted 22 March 2024

ABSTRACT

The paper presents results on tree snails based on samples collected from various parts of Vietnam during the period of 2008–2022. As a result, 14 *Amphidromus* species were recorded, which increased the diversity of the genus in Vietnam to a total of 27 species. Four species were new country records including *Amphidromus atricallosus perakensis*, *Amphidromus leucoxanthus*, *Amphidromus perversus natunensis*, and *Amphidromus areolabiatus*. Almost all 14 species were found in the south, while only *Amphidromus roseolabiatus* was recorded in the north. The great variation in the shell morphology of these tree snails required an integrative taxonomic approach to better understand the species diversity of the genus *Amphidromus* in Vietnam.

Keywords: Mollusca, land snails, checklist, diversity, Vietnam.

Citation: Do Van Nhuong, Dinh Dieu Thuy, Do Duc Sang, Nguyen Thanh Tung, Le Hung Anh, Arthur E. Bogan, Do Van Tu, 2024. Tree snails *Amphidromus* Albers, 1850 (Pulmonata: Camaenidae) collected in Vietnam. *Academia Journal of Biology*, 46(1): 115–133. <https://doi.org/10.15625/2615-9023/18908>

*Corresponding author email: dinhdieuthuy@outlook.com.vn

INTRODUCTION

The tree snail genus *Amphidromus* Albers, 1850 belongs to the family Camaenidae Pilsbry, 1895, and has a wide distribution across northeast India, Myanmar, Malaysia, Thailand, Laos, Cambodia, Vietnam, Indonesia, the southern Philippines, and China (Solem, 1983; Laidlaw & Solem, 1961). The genus *Amphidromus* features an elongated conic or ovate conic helicoid shell comprising six to eight whorls. The shell may vary from thin and fragile to very thick and solid. In some species, the shell consistently coils to the right (dextral), while in many others, it coils consistently to the left (sinistral). Notably, some species exhibit both left- and right-handed coiling within the same species (Laidlaw & Solem, 1961). Many species and varieties were named during the first half of the nineteenth century, often accompanied by inadequate locality data (Laidlaw & Solem, 1961). The taxonomy of this genus has proven intricate due to the diverse shell morphologies and species distribution. In 1896, Fulton organized 142 specific and varietal names into 18 groups, encompassing 64 species. Pilsbry (1900) expanded the number of species by 81, categorized them into 19 groups. Laidlaw & Solem (1961) discussed the variation in *Amphidromus* shell colours across different areas, as well as the factors contributing to changing shell patterns in many species. This monograph has been deemed an important reference for classifying *Amphidromus* based on shell morphology. As of 2015, 95 species have been recognized (Inkhavilay et al., 2017).

Vietnam's wealth of terrestrial biological diversity arises from its diverse geography and climate. The country lies within a humid monsoon region, characterized by numerous mountains, hills, and diverse natural landscapes. The climate, topography, hydrological features, and vegetation in Vietnam collectively contribute to its rich terrestrial biological diversity (Lap, 2009), which includes land snails. Studies on this group in Vietnam commenced in the mid-19th century (Souleyet, 1852; Crosse & Fischer, 1863; Mabille, 1887). Thanh (2008)

documented 448 land snail species in Vietnam. In 2011, Schileyko published a list of 477 pulmonate land snail species and subspecies for the country. The list encompassed 23 *Amphidromus* species. This list is the most comprehensive synthesis of the genus *Amphidromus* in Vietnam. Generally, Schileyko (2011) primarily synthesized the findings of prior studies with limited specimen analysis. Furthermore, distribution information for many species remained unclear.

The current study aims to contribute additional data regarding the diversity of *Amphidromus*. This was achieved through the identification and analysis of samples collected from diverse regions across Vietnam over a period of more than ten years.

MATERIALS AND METHODS

The specimens were collected from over 30 sites across 11 provinces in Vietnam, including both mainland areas and some islands, during the period from 2008 to 2022 (Fig. 1). The mainland sites encompass limestone mountains in Cuc Phuong National Park (Ninh Binh province), Phong Nha-Ke Bang National Park (Quang Binh province), Vinh Cuu (Dong Nai province), Phu Yen, Kien Luong, Ha Tien, and Hon Dat (Kien Giang province). Additionally, some coastal islands were included: Hon Tre, Lai Son, An Son (Kien Giang province), and Con Dao (Ba Ria-Vung Tau province). For some specimens, their shells were cleaned, dried, and stored in plastic boxes. Living specimens were preserved in 80% ethanol. These were photographed using a digital camera to document their coloration. The taxonomic classification system of species along with their synonyms is based on MolluscaBase eds. (2023). The collected specimens have been deposited at the Vietnam National Museum of Nature (VNMN), Hanoi National University of Education (HNUE), and Can Tho University (CTU), and are listed in Table 1.

All tree snail species were identified according to the original descriptions in Fulton (1901), Bavay & Dautzenberg (1909a, b), Laidlaw & Solem (1961), Sutcharit &

Panha (2006), Sutcharit et al. (2013), and Jirapatrasilp et al. (2022). Moreover, we conducted comparisons between our specimens and the illustrated type catalog of *Amphidromus* held at the National History Museum, London (NHMUK), as outlined in Sutcharit et al. (2015).




Figure 1. Sampling locations of *Amphidromus* in Viet Nam

Height (H) measurements were taken from the apex to the base of the shell, in parallel with the shell's axis and perpendicular to the diameter of the shell at its widest point on the

last whorl, including the lip. Width (W) represents the maximum distance between the two sides of the body whorl of the shell. All measurements for adult samples were recorded in millimeters using a micrometer caliper. The values presented for H and W dimensions are the averages derived from measurements taken across all individuals at a given sampling site. Whorls in undamaged adult shells were counted following the methodology of Oesch et al. (2013). All specimens were photographed in standard positions, including the apertural view. We referred to Molluscabase (<https://molluscabase.org/>) for valid names and synonyms. In this context, the abbreviations used include A for adult, J for juvenile, and NP for national park.

RESULTS

The collected specimens were identified as 14 different species and subspecies (Table 1). All *Amphidromus* species recorded in Vietnam belong to two subgenera: *Amphidromus* (*Amphidromus*) Albers, 1850 and *Amphidromus* (*Syndromus*) Pilsbry, 1900. These subdivisions are primarily differentiated by variations in shell size, thickness, colouration, and the morphology of reproductive organs (Laidlaw & Solem, 1961; Sutcharit & Panha, 2006). Five species, *Amphidromus inversus annamiticus*, *Amphidromus cambojiensis*, *Amphidromus cochinchinensis*, *Amphidromus ingens*, and *Amphidromus smithii*, were re-collected from their type localities.

Table 1. The average size of our *Amphidromus* specimens collected in Vietnam
(The name of provinces where the specimens were collected,
and voucher codes were shown below each species)

No.	Species name, locality, and museum code	Number of adults	H (mm)	W (mm)	H/W ratio	Whorl ranges
<i>Amphidromus atricallosus</i> (Gould, 1843)						
1	Dong Nai, HNUE-LS 000.29	3	59.8 ± 2.3	32.0 ± 0.3	1.86	6–7
	Kien Giang, CTU-LS 014.01	2	58.2 ± 0.6	31.3 ± 0.1	1.85	6–7
	Kien Giang, CTU-LS 012.09	3	59.9 ± 3.4	32.0 ± 0.4	1.87	6–7
<i>Amphidromus atricallosus perakensis</i> Fulton, 1901						
2	Kien Giang, HNUE-LS 000.09	2	54.3 ± 0.4	31.5 ± 0.5	1.71	6–6 ^{1/2}
	Kien Giang, HNUE-LS 000.37	3	60.0 ± 0.7	32.4 ± 0.2	1.85	6–6 ^{1/2}

No.	Species name, locality, and museum code	Number of adults	H (mm)	W (mm)	H/W ratio	Whorl ranges
3	Kien Giang, CTU-LS 015.07	3	49.5 ± 0.2	30.2 ± 0.2	1.63	6 ^{1/4}
	Con Dao, VNMN-IZ 000.002.320	1	53.2	29.3	1.81	6
3	<i>Amphidromus cambojiensis</i> (Reeve, 1860)					
	Dong Nai, HNUE-LS 000.01	5	67.6 ± 0.7	35.4 ± 0.8	1.90	6 ^{1/2} –7 ^{1/2}
	Dong Nai, HNUE-LS 000.02	7	65.9 ± 1.9	35.0 ± 0.9	1.87	6 ^{1/2} –7
	Dong Nai, HNUE-LS.000.04	12	66.7 ± 2.6	35.7 ± 2.7	1.86	6 ^{1/2} –7
	Dong Nai, VNMN-IZ 000.002.305	2	64.3 ± 2.4	34.0 ± 0.2	1.89	6 ^{1/2} –7
	Dong Nai, VNMN-IZ 000.002.306	1	70.2	37.0	1.89	6–6 ^{1/2}
4	<i>Amphidromus cochinchinensis</i> (Pfeiffer, 1857)					
	Kien Giang, HNUE-LS 000.08	1	45.0	22.2	2.02	6 ^{1/2}
5	<i>Amphidromus comes</i> (L. Pfeiffer, 1861)					
	Dong Nai, HNUE-LS 000.06	3	50.2 ± 1.7	27.3 ± 0.9	1.84	6 ^{1/4} –7
	Dong Nai, HNUE-LS 000.07	17	47.0 ± 2.2	26.1 ± 2.1	1.79	6–6 ^{3/4}
	Kien Giang, VNMN-IZ 000.002.319	1	46.4	25.1	1.84	6 ^{1/4}
	Dong Nai, VNMN-IZ 000.002.307	3	44.4 ± 0.2	27.4 ± 0.3	1.62	6–6 ^{1/2}
6	<i>Amphidromus ingens</i> Möllendorff, 1900					
	Phu Yen, VNMN-IZ 000.002.318	3	60.4 ± 0.4	33.9 ± 0.3	1.78	6–6 ^{1/2}
7	<i>Amphidromus inversus annamiticus</i> (Crosse et Fischer, 1863)					
	Kien Giang, HNUE-LS 000.30	4	48.6 ± 2.7	26.0 ± 0.9	1.86	6 ^{3/4}
	Kien Giang, CTU-LS 012.01	6	51.8 ± 2.3	30.2 ± 1.1	1.71	6 ^{1/2}
	Kien Giang, CTU-LS 012.03	10	48.7 ± 2.4	25.5 ± 2.1	1.90	6 ^{1/2}
	Kien Giang, CTU-LS 012.04	4	49.6 ± 0.7	27.2 ± 0.6	1.82	6 ^{1/2}
	Kien Giang, CTU-LS 012.05	6	50.2 ± 1	28.1 ± 0.4	1.78	6–6 ^{1/4}
8	<i>Amphidromus leucoxanthus</i> (von Martens, 1864)					
	Kien Giang, HNUE-LS 000.36	1	57.8	31.5	1.77	6 ^{3/4} –7
	Kien Giang, CTU-LS 015.01	7	51.2 ± 2.4	26.4 ± 1.4	1.93	6 ^{1/2}
	Kien Giang, CTU-LS 015.02	13	52.6 ± 1	28.4 ± 0.7	1.84	6 ^{1/4} –6 ^{3/4}
	Kien Giang, CTU-LS 015.04	11	52.3 ± 1.6	27.2 ± 0.3	1.92	6 ^{1/2}
	Kien Giang, CTU-LS 015.05	15	52.1 ± 1.2	27.6 ± 0.5	1.88	6 ^{1/2}
9	<i>Amphidromus mouhoti</i> (Pfeiffer, 1861)					
	Dong Nai, VNMN-IZ 000.002.316	1	34.3	16.5	2.07	6 ^{1/4}
	Quang Binh, VNMN-IZ 000.002.317	2	35.9 ± 0.3	20.2 ± 0.1	1.77	6
10	<i>Amphidromus perversus natunensis</i> Fulton, 1906					
	Kien Giang, CTU-LS 015.11	4	49.1 ± 1.4	29.0 ± 1.2	1.68	6–6 ^{1/4}
	Kien Giang, CTU-LS 015.12	5	50.3 ± 1.3	28.3 ± 2.1	1.77	6–6 ^{1/2}
	Con Dao Island, VNMN-IZ 000.002.309	1	58.7	30.0	1.95	6 ^{1/2}
11	<i>Amphidromus roseolabiatus</i> Fulton, 1896					
	Quang Binh, VNMN-IZ 000.002.314	4	37 ± 0.4	20.6 ± 0.1	1.79	6 ^{1/4}
12	<i>Amphidromus areolatus</i> (L. Pfeiffer, 1861)					
	Tay Ninh, HNUE-LS 000.10	1	25.5	12.3	2.07	5 ^{1/2}
13	<i>Amphidromus smithii</i> Fulton, 1896					
	Dong Nai, HNUE-LS 000.12	1	35.5	15.6	2.27	6 ^{1/2}
	Dong Nai, VNMN-IZ 000.002.311	1	32.5	14.2	2.28	6 ^{1/4}
	Dong Nai, VNMN-IZ 000.002.312	1	36.0	17.2	2.09	6
	Dong Nai, VNMN-IZ 000.002.313	1	35.0	15.4	2.27	6
14	<i>Amphidromus zebrinus</i> (Pfeiffer, 1861)					
	Dong Nai, HNUE-LS 000.11	2	23.0 ± 0.3	12.1 ± 0.1	1.90	6

Checklist of *Amphidromus* species in Vietnam (*collected in this study; **from checklist by Schileyko (2011) and not obtained in this study)

Family Camaenidae Pilsbry, 1895

Subfamily Camaeninae Pilsbry, 1895

Genus *Amphidromus* Albers, 1850

Subgenus *Amphidromus* Albers, 1850

Bulimus (*Amphidromus*) Albers, 1850: 138.

Type species. *Helix perversus* Linnaeus, 1758, by the subsequent designation of von Martens (1860: 184).

Amphidromus asper* Haas, 1934*

Type locality. "Süd-Annam, 120 km von der Küste, auf dem Wege zum Plateau von Lang-Bian, zw. 600-1000 H" (Central Vietnam, 120 km on the way from the coast to the Lang Bian Plateau, 600-1000 meters elevation).

Amphidromus atricallosus* (Gould, 1843)
(Fig. 2a)

Bulimus atricallosus Gould, 1843: 142.

Bulimus eques Pfeiffer, 1857: 158;
Fischer, 1891: 30, 31.

Amphidromus (*Amphidromus*) *atricallosus*
Laidlaw & Solem, 1961: 530; Sutcharit &
Panha, 2006: 14.

Amphidromus atricallosus Schileyko,
2011: 49.

Type locality. Tavoy, Burmah.

Material examined. VIETNAM: Dong Nai province: 3A, HNUE-LS 000.29, Vinh Cuu district, Dong Nai Culture and Nature Reserve, 2008, Nhuong DV leg.; Kien Giang province: 2A, CTU-LS 014.01, Kien Hai district, Lai Son Island, natural forest, 9°47'55.1"N - 104°37'22.9"E, alt. 33 m, 14.xi.2013, Be NV leg.; Kien Giang province: 3A, CTU-LS 012.09, Hon Dat district, 10°09'53.0"N - 104°53'53.0"E, vii.2008, Be NV leg.

Distribution. Vietnam (Dong Nai, Kien Giang, Ninh Thuan, Khanh Hoa), Malaysia,

Lower Myanmar, South of Thailand (Schileyko, 2011).

Remarks. The materials have the same characters as *Bulimus atricallosus* (Gould, 1843) such as a solid shell, imperforate conical shape, with 6-7 slightly convex whorls, lunate-oval aperture, white and reflected lip; they also match with the description in Laidlaw & Solem (1961): substantial size and have white sub-band below the suture. Variability in our specimens is observed in both sinistral and dextral shells, displaying coloration that spans from yellow green to light yellow or white; the apex tends to be light yellow to white; the axis may be either straight or curved, and the parietal is slightly black or milky white; some individuals exhibit a slit-like umbilicus. Notably, this species presents various morphological variations across different regions (Sutcharit & Panha, 2006), suggesting the need for further research.

Amphidromus atricallosus perakensis* Fulton, 1901 (Fig. 2b)

Amphidromus perakensis Fulton, 1901: 104, pl. 9, figs. 8-10.

Amphidromus (*Amphidromus*) *atricallosus*
perakensis Sutcharit & Panha, 2006: 21,
figs. 4F, G.

Type locality. Perak (Peninsular Malaysia).

Material examined. VIETNAM: Kien Giang province: 3A, 2J, CTU-LS 015.07, Kien Luong district, Binh An limestone hill, 10°09'49.5"N - 104°37'12.1"E, 8.xii.2014, Be Hai VV leg.; Kien Giang province: 2A, CTU-LS 015.08, Ha Tien city, Thach Dong limestone hill, 10°24'39.4"N - 104°28'29.4"E, 9.xii.2014, Be Hai VV leg.; Kien Giang province: 1A, CTU-LS 015.09, Kien Hai district, Hon Tre Island, 09°58'19.0"N - 104°50'49.5"E, 21.x.2020, Be NV leg.; Kien Giang province: 3A, 5T, CTU-LS 015.10, Kien Hai district, Hon Tre island, 09°58'19.5"N - 104°50'46.5"E, 14.xi.2013, Be NV leg.; Kien Giang province: 2A, 2J, HNUE-LS 000.09, Kien Hai district, Hon Tre Island, 09°58'19.3"N - 104°50'46.5"E,

21.x.2020, Nhuong DV leg.; Kien Giang province: 3A, HNUE-LS 000.37, Hon Dat district, $10^{\circ}09'53.0''$ N - $104^{\circ}53'53.0''$ E, vii.2008, Be NV leg.; Ba Ria-Vung Tau province: 1A, VNMN-IZ 000.002.320, Con Dao Island, $8^{\circ}42'31.313''$ N - $106^{\circ}35'49.445''$ E, 18.v.2022, Tu DV leg.

Distribution. Vietnam (Kien Giang, Con Dao island), southern Malaysia peninsula (including Singapore) (Sucharit & Panha, 2006).

Remarks. Our specimens were identified as *Amphidromus atricallosus perakensis* by having some characteristics such as there are both dextral and sinistral shells, yellow shell with a conspicuous band at the suture, slightly convex, peristome white, thick, expanded and slightly reflected (Fulton, 1901). Specimens collected in Vietnam show variation in size (see Table 1).

Amphidromus cambojiensis* (Reeve, 1860)
(Fig. 2c)

Bulimus cambojiensis Reeve, 1860: 204.

Amphidromus cambojiensis—Laidlaw & Solem, 1961: 529; Schileyko, 2011: 50; Sutcharit et al., 2015: 62, figs. 4F–G.

Amphidromus lamdongensis Thach & Huber in Thach, 2016: 67, figs. 32, 339–342.

Amphidromus schileykoi Thach, 2016: 68, figs. 39, 381–383.

Amphidromus montesdeocai Thach & Huber in Thach, 2017: 43, figs. 454–458.

Type locality. Cambodia (Cambodia).

Material examined. VIETNAM: Dong Nai province: 8A, 1J, HNUE-LS 000.01, Vinh Cuu district, forest in Dakinde, $11^{\circ}10'18.03''$ N - $107^{\circ}02'06.88''$ E, 24.v.2010, Nhuong DV leg.; Dong Nai province: 7A, HNUE-LS 000.02, Vinh Cuu district, forest around Ba Hao lake, $11^{\circ}16'40.26''$ N - $107^{\circ}05'12.13''$ E, 23.v.2010, Nhuong DV leg.; Dong Nai province: 1J, HNUE-LS 000.03, Vinh Cuu district, forest in zone No. 93, $11^{\circ}16'19.39''$ N - $107^{\circ}04'34.24''$ E, 22.v.2010, Nhuong DV leg.; Dong Nai province: 12A, 1J, HNUE-LS 000.04, Tan Phu district, the

forest in Cat Tien NP near Bau Chim, $11^{\circ}29'13.67''$ N $107^{\circ}22'55.05''$ E, 20.v.2021, Nhuong DV leg.; Dong Nai province: 12A, HNUE-LS 000.05, Vinh Cuu district, humid forest, $11^{\circ}14'38.13''$ N - $107^{\circ}03'34.13''$ E, Hung NV leg., v.2021; Kien Giang province: 1A, CTU-LS 013.02, Hon Dat district, Hon Me mountain, natural forest, $10^{\circ}06'30.6''$ N - $104^{\circ}53'23.7''$ E, alt. 83 m, 07.xii.2014, Be Hai VV leg.; Kien Giang province: 1A, CTU-LS 013.03, Ha Tien city, Thach Dong mountain, natural forest, $10^{\circ}24'39.4''$ N - $104^{\circ}28'29.4''$ E, alt. 25 m, 09.xii.2014, Be Hai VV leg.; Kien Giang province: 1A, CTU-LS 013.05, Hon Dat district, Hon Soc mountain, natural forest, $10^{\circ}09'25.3''$ N - $104^{\circ}54'14.2''$ E, alt. 60 m, 07.xii.2014, Be Hai VV leg.; Dong Nai province: 2A, VNMN-IZ 000.002.305, Vinh Cuu district, Hieu Liem commune, forest in Da Dung, $11^{\circ}11'1.051''$ N - $106^{\circ}57'32.871''$ E, 9.vi.2020, Thuy DD leg.; Dong Nai province: 1A, VNMN-IZ 000.002.306, Vinh Cuu district, Phu Ly commune, Bu Dang forest, $11^{\circ}26'27.5''$ N - $107^{\circ}06'08.6''$ E, 11.vii.2020, Thuy DD leg.

Distribution. Vietnam (Dong Nai, Kien Giang, Lam Dong) (Pall-Gergely et al., 2020), Cambodia (Schileyko, 2011).

Remarks. *Amphidromus cambojiensis* is the largest species in this genus (see Table 1). Our materials share some key characters with the original description of *Bulimus cambojiensis* by Reeve (1860) such as: shell dextral or sinistral; whorls $6-7^{1/2}$, smooth, convex; ovate aperture, with white, thick and reflected lip. However, a comparative analysis between our specimens and the original description by Reeve (1860) reveals some distinctions: there is a variation in shell colors, ranging from deep brown to violet-rose, with certain individuals displaying a milky-white or light grey interior; and slightly smaller (H 61.9–70.2 mm vs. 76 mm).

The type locality of this species remains unclarified. In 1860, Reeve described this species from Mouhot's specimens collected in the Kingdom of Cambodia without giving the exact data of the type locality. Laidlaw & Solem (1961) and Schileyko (2011) stated

that the species had been described from “60 leagues north of Saigon, in the Stiengs country or Moi”, which is in Vietnam. Therefore, further research is needed to determine the current type of locality in which country.

Amphidromus chloris* (Reeve, 1848)*

Type locality. “Eastern Islands” (West Australia).

***Amphidromus cochinchinensis* (L. Pfeiffer, 1857)* (Fig. 2d)**

Bulimus cochinchinensis Pfeiffer, 1857 [1856]: 331, 332.

Amphidromus metabletus Möllendorff, 1900; Laidlaw & Solem, 1961: 528.

Amphidromus metabletus insularis Möllendorff, 1901.

Amphidromus metabletus pachychilus Möllendorff, 1901.

Amphidromus cochinchinensis Schileyko, 2011: 50; Sutcharit et al., 2015: 64, figs. 5D–E.

Type locality. Cochin China (Southern Vietnam).

Material examined. VIETNAM: Kien Giang province: 1A, HNUE-LS 000.08, Hon Dat district, Tho Son commune, Hon Me mountain, 10°06'30.6"N - 104°53'23.7"E, 7.xii.2014, Be NV leg.

Distribution. Vietnam (Khanh Hoa, Kien Giang) (Laidlaw & Solem, 1961)

Remarks. Compared with the original description in Pfeiffer (1857), the remarks of Laidlaw & Solem (1961), and figs. 5D-E by Sutcharit et al. (2015), our specimens are identified as *Amphidromus cochinchinensis*. They share some main features including: Shell shape (conical elongated), the aperture is oblique, sublunate, and moderately thick with a reflected lip; the axis is straight. However, our specimen is larger in size (45 mm vs. 39mm) (Laidlaw & Solem, 1961) and difference in color (yellow vs white) (Sutcharit et al., 2015).

***Amphidromus comes* (L. Pfeiffer, 1861)* (Fig. 2e)**

Bulimus comes L. Pfeiffer, 1861: 193.

Cochlostyla polymorpha Tapparone-Canefri, 1874: 82, plate 2, figs. 4a–b.

Amphidromus comes polymorphus Pilsbry, 1900: 171, plate 57, figs 6–10.

Amphidromus comes-Pilsbry 1900: 170, plate 57, figs. 1–5; Laidlaw & Solem, 1961: 531; Schileyko, 2011: 50; Sutcharit et al., 2015: 64, Fig. 5 G-H; Inkhavilay et al., 2019: 89; Pall-Gergely et al., 2020: 51.

Amphidromus hueae Thach & Huber, 2016 in Thach 2016: 66, figs 38, 331–334.

Amphidromus ngocngai Thach, 2017: 44, figs. 446–449.

Amphidromus vietnamensis Thach & Huber, 2016 in Thach 2017: 48, figs. 630–635.

Amphidromus dongnaiensis Thach, 2018: 51, 789–793.

Amphidromus atricallosus vovanae Thach, 2019: 84.

Type locality. Cambodia.

Material examined. VIETNAM: Dong Nai province: 8A, 22J, HNUE-LS 000.06, Vinh Cuu district, natural forest, 11°14'38.13"N - 107°03'34.13"E, 25.v.2021, Hung NV leg.; Dong Nai province: 17A, 40T, HNUE-LS 000.07, Tan Phu district, a natural forest in Cat Tien NP, 11°27'44.56"N - 107°19'16.98"E, Nhuong DV leg., 20.v.2021; Dong Nai province: 3A, VNMN-IZ 000.002.307, Long Khanh district, Suoi Tre commune, 6.vii.2019, Tien BT leg.; Dong Nai Province: 2A, VNMN-IZ 000.002.308, Vinh Cuu district, Ma Da commune, the park near Ba Hao lake, 11°15'32.25"N - 107°4'19.774"E, 12.vi.2020, Thuy DD leg.; Kien Giang province: 1A, VNMN-IZ 000.002.319, Tho Chu Island, 21.vi.2022, Tu D.V leg.

Distribution. Vietnam (Dong Nai, Kien Giang), Laos, Thailand, Cambodia (Laidlaw & Solem, 1961; Schileyko, 2011).

Remarks. Our materials share some main characters with *Amphidromus comes* such as:

shell dextral or sinistral, solid, conical spire, oblique aperture (Pilsbry, 1900), the apex is dark (Laidlaw & Solem, 1961) (Sutcharit et al., 2015, figs. 5G–H). Nonetheless, specimens collected from Vietnam exhibit some

distinctions, including variations in shell color (including unpolished shades of yellow, brown-purple, and grayish-green), as well as uneven embryonic and post-embryonic whorls in some individuals, resembling malformations.



Figure 2. a. Amphidromus atricallosus, CTU-LS 012.09; b. Amphidromus atricallosus perakensis, CTU-LS 015.10; c. Amphidromus cambojiensis, HNUE-LS 000.05; d. Amphidromus cochinchinensis, HNUE-LS 000.08; e. Amphidromus comes, HNUE-LS 000.07; f. Amphidromus ingens, VNMN-IZ 000.002.318. Scale: 10 mm

Amphidromus costifer* Smith, 1893*

Type locality. "Montagnes boîties du Huyen de Tri-phuoc, Province Binh-dinh, Annam" (Binh Dinh province, Vietnam).

Amphidromus dautzenbergi* Fulton, 1899*

Type locality. Tonkin.

Amphidromus dohrni* (Pfeiffer, 1863)*

Type locality. Cochin-China.

Amphidromus ingens* Möllendorff, 1900
(Fig. 2f)

Amphidromus ingens Möllendorff, 1900a: 23; Zilch, 1953: 135, plate 23, fig. 25; Schileyko, 2011: 50.

Amphidromus naggisi Thach & Huber, 2014: 35, figs. 1–13, 15.

Type locality. "Berg Mutter und Kind" (Mother and Child Mountain - Central Vietnam).

Material examined. VIETNAM: Phu Yen province: 3A, VNMN-IJ 000.002.318, Tuy Hoa district, the forest in Hoa My Dong, 12°50'00"N - 109°12'00.9"E, 7.iv.2022, Thuy DD leg.

Distribution. Vietnam (Phu Yen, Mother and Child Mountain - Central Vietnam) (Schileyko, 2011).

Remarks. Our specimens display the main characteristics of this species, such as a thick shell with rough surface due to the grooves spread from the last whorl up to the near apex, thick lip and grey or milky nacre layer. The shell showed both dextral and sinistral coiling patterns (Möllendorff, 1900). Compared with the image of Zilch (1953), the shell shape is similar.

***Amphidromus inversus annamiticus* (Crosse & Fischer, 1863)* (Fig. 3a)**

Bulimus annamiticus Crosse & Fischer, 1863: 357–359; Crosse & Fischer, 1864: 329, fig 8, pl. XII; Mabille & Mesle, 1866: 128.

Amphidromus annamiticus—Morlet, 1889: 126.

Amphidromus annamiticus var. *roseotincta* Möllendorff, 1894: 150.

Amphidromus inversus annamiticus—Pilsbry, 1900: 169–170, pl. 56, figs 98–100; Laidlaw & Solem, 1961: 561, 600; Solem, 1965: 624; Schileyko, 2011: 50; Sutcharit et al., 2013: 53–57, figs. 1 A–D; Inkhvilay et al., 2019: 91, figs. 44 D, E.

Amphidromus inversus annamiticus var. *roseotincta*—Pilsbry, 1900: 170; Laidlaw & Solem, 1961: 561, 655–656; Solem, 1965: 624.

Amphidromus inversus var. *annamiticus*—Gude, 1903a: 50, 51; Abbott, 1989: 160.

Amphidromus inversus roseotincta—Zilch, 1953: 135, pl. 23, fig. 26.

Amphidromus (Amphidromus) inversus annamiticus Sutcharit & Panha, 2006: 9, figs. 3E–H.

Type locality. "in vicinio urbis Saigon et pagi Fuyen-Moth dicti" (Ho Chi Minh city).

Material examined. VIETNAM: Kien Giang province: 6A, CTU-LS 012.01, Kien Hai district, Hon Tre Island, permanent cropland, 09°58'30.5"N - 104°51'23.0"E, alt. 41 m, Be NV leg., 30.x.2013; Kien Giang province: 10A, CTU-LS 012.03, Kien Luong district, Binh An mountain, permanent cropland of *Aquilaria* sp., 10°09'49.5"N - 104°37'12.1"E, alt. 22 m, 8.xii.2014, Be Hai VV leg.; Kien Giang province: 4A, CTU-LS 012.04, Kien Hai district, Lai Son Island, permanent cropland, 09°48'10.7"N - 104°37'53.6"E, alt. 27 m, 14.x.2014, Be NV leg.; Kien Giang province: 6A, CTU-LS 012.05, Kien Hai district, An Son Island, land for short-term crops, 09°40'42.3"N - 104°21'10.8"E, alt. 270 m, Be NV leg., 17.xi.2013; An Giang province: 5A, CTU-LS 012.06, Thoai Son district, Ba The mountain, 9.ix..2014, Be Hai VV leg.; Kien Giang province: 4A, HNUE-LS 000.30, Hon Dat district, Hon Soc village, 10°09'53.0"N - 104°53'53.0"E, 8.xii.2014, Be Hai VV leg.

Distribution. Vietnam (Kien Giang, An Giang), Thailand (in the South including some islands (Laidlaw & Solem, 1961; Sutcharit, 2013), Campuchia (Lower Mekong Valley) (Pilsbry, 1900).

Remarks. The specimens are identified as *Amphidromus inversus annamiticus*, sharing

some main characteristics described by Crosse & Fischer (1863) including ovate-conical shell, dark brown stripes, spirally banded with a dark purple stripe, conical spire, slightly obtuse apex, violet-blackish. When contrasted with specimens from Thailand (Sutcharit & Panha, 2006), ours exhibit distinct differences. In Vietnam, sinistral shells are significantly more common (making up about 75%); shell sizes are larger (ranging from H 48.6–51.8mm and W 26–30.2mm), whereas in Thailand, most specimens have a dextral helix; and the shell sizes range from H 32.7–51.4mm and W 19.8–28.2mm. Furthermore, the apex color in Vietnam specimens is limited to pink or black, while in Thailand (figs. 3E–H, cf. Sutcharit & Panha, 2006), it encompasses a wider spectrum of colors such as pink, ivory, purple, or white in some cases.

Amphidromus mirandus* Bavay et Dautzenberg, 1912*

Type locality. Lang-Biang, Annam

***Amphidromus leucoxanthus* (von Martens, 1864)* (Fig. 3b)**

Bulimus leucoxanthus von Martens, 1864: 526; von Martens, 1867: 348–349, pl. 20, figs. 11,12.

Amphidromus leucoxanthus—Morlet, 1889: 127; Laidlaw & Solem, 1961: 530.

Amphidromus perversus var. *leucoxanthus* Fulton, 1896: 69; Laidlaw & Solem, 1961: 635–636

Amphidromus aureus *leucoxanthus* Pilsbry, 1900: 163–164, pl. 54, figs 73–79.

Amphidromus aureus var. *leucoxanthus* Gude, 1903: 8.

Amphidromus atricallosus var. *leucoxanthus* Laidlaw & Solem, 1961: 635–636; Solem, 1965: 620, pl. 1, fig. 8; Abbott, 1989: 160.

Amphidromus (Amphidromus) atricallosus leucoxanthus Sutcharit & Panha, 2006: 20, fig. 3P.

Type locality. Siam, location of types unknown.

Material examined. VIETNAM: Kien Giang province: 7A, CTU-LS 015.01, Kien Hai district, Lai Son Island, permanent cropland, 09°47'55.1"N - 104°37'22.9"E, alt. 33 m, 14.xi.2014, Be NV leg.; Kien Giang province: 13A, CTU-LS 015.02, Ha Tien city, Thach Dong limestone hill, inside the cave, 10°24'39.4"N - 104°28'29.4"E, alt. 23 m, 09.ix.2014, Be Hai VV leg.; Kien Giang province: 1A, CTU-LS 015.03, Hon Dat district, Hon Soc mountain, natural forest, 10°09'25.3"N - 104°54'14.2"E, alt. 60 m, 07.xii.2014, Be Hai VV leg.; Kien Giang province: 11A, CTU-LS 015.04, Kien Hai district, An Son Island, natural forest, 09°41'24.3"N - 104°21'22.9"E, alt. 50 m, 13.x.2014, Be NV leg.; Kien Giang province: 15A, CTU-LS 015.05, Kien Luong district, Binh An mountain, permanent cropland, alt. 22 m, 10°09'49.5"N - 104°37'12.1"E, 08.xii.2014, Be Hai VV leg.; Kien Giang province: 1A, CTU-LS 015.06, Kien Hai district, Hon Tre Island, permanent cropland, 09°58'30.5"N - 104°51'23.0"E, alt. 41 m, 30.x.2013, Be NV leg.; Kien Giang province: 1A, HNUE-LS.000 36, An Son district, Hon Tre island, 21.x.2020, Tung NT leg.

Distribution. Vietnam (Kien Giang), Eastern Thailand, and maybe Cambodia (Laidlaw & Solem, 1961). According to Sucharit & Panha (2006), this species is distributed in the Chachoengsao, Chonburi and Chantaburi provinces of Thailand.

Remarks. Our specimens agree with descriptions of *Bulimus leucoxanthus* by Martens (1864) such as conical-elongated shell, lightly striated, glossy, citrine, apex white, sutura appressed, aperture rounded, thickened and white parietal callus. We found both dextral and sinistral shells of this species in Vietnam. In contrast to the specimens in Thailand, our specimens lack a white band under the suture (compared to the presence of the white band under the suture observed in samples from Chonburi, cf. Sutcharit et al., 2006).

Amphidromus mouhoti* (Pfeiffer, 1861)
(Fig. 3c)

Bulimus mouhoti Pfeiffer, 1861: 194;
Fischer, 1891: 32; Fulton, 1868: 88; Laidlaw
& Solem, 1961: 524.

Amphidromus mouhoti—Schileyko, 2011:
51; Sutcharit et al., 2015: 82, fig 11E; Barna
Pál-Gergely, 2020: 53.

Type locality. Siam (Thailand).

Material examined. VIETNAM: Dong Nai province: 1A, VNMN-IZ 000.002.315, Long Khanh district, Suoi Tre commune, shrubs in the fruit garden, vii.2019, Tien BT leg.; Dong Nai province: 1A, VNMN-IZ 000.002.316, Vinh Cuu district, Ma Da commune, forest near Ba Hao lake, 11°15'43.71"N - 107°05'26.91"E, vi.2020, Thuy DD leg.; Quang Binh province: 2A, VNMN-IZ 000.002.317, Phong Nha - Ke Bang NP, entrance to En Cave, 17°26'03.7"N - 106°17'58.3"E, alt. 214 m, 30.v.2022, Thuy DD leg.

Distribution. Vietnam (Quang Binh, Dong Nai), Thailand, Cambodia (Schileyko, 2011).

Remarks. Pfeiffer (1861) did not include illustrations in the species description, making identification challenging. Nonetheless, cross-referencing with the figure of this species in NHMUK (Sutcharit et al., 2015) may aid in identification, despite potential differences in size. Specimens from Vietnam share the main characteristics of this species such as flat or slightly swollen whorls, a sublunate aperture, purplish or pink lips, a pointed apex, and a light-yellow band parallel to the suture (Pfeiffer, 1861). However, our specimens have variation in coloration, ranging from light yellow transitioning to white at the apex. Notably, Phong Nha-Ke Bang specimens lack a green groove in the last whorl, while Dong Nai specimens exhibit green or grey grooves.

Amphidromus perversus* (Linnaeus, 1758)*

Type locality. unknown

Amphidromus perversus natunensis* Fulton, 1896 (Fig. 3d)

Amphidromus perversus natunensis Fulton, 1896: 69; Laidlaw & Solem, 1961:

536–537, 643, fig. 19; Sutcharit et al., 2015: 82, figs. 11H–I.

Amphidromus aureus natunensis Pilsbry, 1900: 162–163, pl. 55, figs. 87–90.

Amphidromus (Amphidromus) perversus natunensis—Sutcharit & Panha, 2006: 6, figs. 3A, B.

Type locality. Natuna Island, Indonesia.

Material examined. VIETNAM: Kien Giang province, 5A, CTU-LS 015.11, Kien Hai district, Lai Son Island, 09°47'55.0"N - 104°37'22.9"E, 14.xi.2013, Be Hai VV leg.; Kien Giang province, 17A, CTU-LS 015.12, Kien Luong district, Binh An limestone hill, 10°09'49.5"N - 104°37'12.1"E, 8.xii.2014, Be NV leg.; Ba Ria-Vung Tau province, 1A, VNMN-IZ 000.002.309, Con Dao island, 8°42'31.313"N - 106°35'49.445"E, 18.v.2022, Tu DV leg.

Distribution. Vietnam (Kien Giang, Con Dao island), Indonesia (Great Natuna Islands) (Fulton, 1896; Pilsbry 1900; Laidlaw & Solem, 1961).

Remarks. Our materials have some main feature of *Amphidromus perversus natunensis* such as a large shell, depressed at the sutural area (Fulton, 1896); whorls quite flattened; the apex acute, whitish; aperture broad, ovate; peristome white, slightly thickened, folded, expanded (Sutcharit et al., 2015: figs. 11H–I). Nonetheless, the shell coloration in our specimens is diverse, showcasing alternating brown and yellow stripes. This color pattern is observed only from the 4th whorl to the last. As we move from the 4th whorl towards the apex, the predominant hues are light yellow or white. The brown band beneath the suture is discontinuous and frequently interrupted.

Amphidromus placostylus* Möllendorff, 1900*

Type locality. “Phuc-son” (Central Vietnam).

Amphidromus rhodostylus* Möllendorff, 1901*

Type locality. “Pharang, Süd-Annam” (Phan Rang, Central Vietnam).

Amphidromus roseolabiatus* Fulton, 1896
(Fig. 3e)

Amphidromus roseolabiatus Fulton, 1896:
89; Pilsbry, 1900: 188, pl. 60, fig 36; Laidlaw

& Solem, 1961: 527; Schileyko, 2011: 51;
Sutcharit et al., 2015: 88, figs. 13j–k;
Inkhavilay et al., 2017: 3–10; Inkhavilay et
al., 2019: 94, fig 45D-F, 58A; Páll-Gergely et
al., 2020: 54.



Figure 3. a. *Amphidromus inversus annamiticus*, CTU-LS 012.01; b. *Amphidromus leucoxanthus*, CTU-LS 015.02; c. *Amphidromus mouhoti*, VNMN-IZ 000.002.316; d. *Amphidromus perversus natunensis*, CTU-LS 015.02; e. *Amphidromus roseolabiatus*, VNMN-IZ 000.002.314; f. *Amphidromus areolatus*, HNUE-LS 000.10. Scale 10 mm

Type locality. Siam (Thailand).

Material examined. VIETNAM: Quang Binh province, 8A, VNMN-IZ 000.002.314, Phong Nha - Ke Bang NP, entrance to En Cave, alt. 214 m, 17°26'03.7"N - 106°17'58.3"E, 30.v.2022, Thuy DD leg.; Ninh Binh province, 1A, ZMB/Moll 269876, Cuc Phuong NP, the way near Mac Lake, 20°15'13.7"N - 105°42'37.4"E, 04.v.2019, Sulikowska-Drozd and D., Thuy DD leg.

Distribution. Vietnam (Phong Nha - Ke Bang NP, Cuc Phuong NP), Laos (Khammouan, Bolikhampasai, Luang Phrabang, and Vientiane), Thailand (Inkhavilay et al., 2017).

Remarks. Our samples were identified as *Amphidromus roseolabiatus* based on some main characteristics including ovate-conic shell, moderately umbilicated, whorls slightly convex, the lip and columella are pink and expanded, with the lip somewhat reflected, while the interior of the aperture is white (Fulton, 1896). Compared with Lao's specimens, our materials have some differences: The size of our specimens are approximately 37 mm in height and 20.6 mm in width, while the shell dimension in Laos varies greatly (H 26.6–37.5 mm, W 12.9–20.2 mm) (Inkhavilay et al., 2017); in addition, the colour of our shell surface is not green as the specimens in Laos (Sutcharit et al., 2015, figs. 13j–k; Inkavilay et al., 2019, figs. 45D–F), it is light yellow.

Subgenus *Syndromus* Pilsbry, 1900

Amphidromus (Syndromus) Pilsbry, 1900: 184.

Type species. *Helix contraria* Müller, 1774 by subsequent designation in Zilch (1960: 623).

Amphidromus areolatus* Pfeiffer, 1861
(Fig. 3f)

Bulimus areolatus L. Pfeiffer, 1861: 194.

Amphidromus areolatus—Sutcharit et al., 2015: 58, fig. 3j–k; Inkavilay et al., 2019: 88, fig. 42, D; Pál-Gergely et al., 2020: 49.

Amphidromus (Syndromus) areolatus
Inkhavilay et al., 2017: 28, fig. 10C.

Type locality. Siam [Thailand].

Material examined. VIETNAM: Tay Ninh province, 1A, HNUE-LS 000.10, Tan Bien district, Tan Lap commune, 11°44'02.14"N - 106°03'07.84"E, 25.vii.2010, Nhuong DV leg.

Distribution. Vietnam (Tay Ninh), Laos, Thailand (Sutcharit et al., 2017).

Remarks. Our specimen agrees with the description of *Bulimus areolatus* by Pfeiffer (1861) including thin and ovate-conical shell; brown conical spire; whorls slightly convex; columella slightly twisted; aperture oblique; peristome thin, slightly expanded.

Amphidromus flavus* (Pfeiffer, 1861)*

Type locality. Siam (Thailand).

Amphidromus fultoni* (Ancey, 1897)*

Type locality. Cochinchina.

Amphidromus semitessellatus* (Morlet, 1885)*

Type locality. "Les montagnes qui bordent le grand fleuve au delà de Stung-Treng. Les forêts et les montagnes de Kampot à Compong-Som" (Cambodia)

Amphidromus smithii* Fulton, 1896 (Fig. 4a)

Amphidromus smithii—Fulton, 1896: 88, Pl. VII, figs. 12, 12a; Möllendorff, 1900: 132; Schileyko, 2011: 51; Sutcharit et al., 2015: 91, figs. 14 I, J; Pál-Gergely et al., 2020: 55.

Amphidromus smithii ventrosulus
Möllendorff, 1900: 132.

Amphidromus noriokowasoei Thach & Huber, 2017a.

Amphidromus tedbaeri Thach, 2017b: 37, figs. 18–20.

Amphidromus baerorum Thach, 2017c: 297, figs. 6–9.

Amphidromus christabaerae Thach, 2017c: 296, figs. 1–4.

Amphidromus steveni Thach, 2017b: 37, figs. 18–20.

Amphidromus eboricolour Thach, 2018a: 51, figs. 795–798.

Amphidromus davidmonsecouri Thach, 2018a: 50, figs. 803–807.



Figure 4. a. *Amphidromus smithii*, VNMN-IJ 000.002.311; b. *Amphidromus zebrinus*, HNUE-LS 000.11. Scale 10 mm

Amphidromus gittenbergeri Thach & Huber, 2018a: 53, figs. 670–675.

Amphidromus noriokowasoei Thach, 2018a: figs. 810–811.

Amphidromus semicinereus Thach, 2018a: 62, figs. 603–606.

Type locality. Phuc Son, Annam (Central Vietnam).

Material examined. VIETNAM: Dong Nai province, 1A, 3J, HNUE-LS 000.12, Vinh Cuu district, forest in Sub-area No.93 in Dong Nai Culture and Nature Reserve, 11°16'19.39"N - 107°04'34.24"E, 22.v.2010, Nhuong DV leg.; Dong Nai province, 1A, 1J, VNMN-IJ 000.002.310, Vinh Cuu district, Ma Da commune, forest near Ba Hao lake, 11°15'43.71"N - 107°05'26.91"E, vi.2020, Thuy DD leg.; Dong Nai province, 1A, VNMN-IJ 000.002.311, Vinh Cuu district, Hieu Liem commune, forest in Da Dung, 11°11'1.051"N - 106°57'32.871"E, 9.vi.2020, Thuy DD leg.; Dong Nai province, 1A, VNMN-IJ 000.002.312, Vinh Cuu district, Phu Ly commune, 11°23'3.932"N - 107°3'42.178"E, 8.vi.2020, Thuy DD leg.; Dong Nai province, 1A, VNMN-IJ 000.002.313, Vinh Cuu district, Phu Ly commune, Bu Dang forest,

11°26'27.5"N - 107°06'08.6"E, 11.vi.2020, Thuy DD leg.

Distribution. Vietnam (Dong Nai).

Remarks. The examined specimens have a weathered stratum corneum, such that the patterns on the shell are not clear. However, they appear like the type specimens shown in Sutcharit et al. (2015) in shell shape, aperture, and lip color. Especially, the periostracum of the shell is very thin and easy to peel off.

Amphidromus ventrosulus Möllendorff, 1900**

Type locality. "Phuc-son, Annam" (Central Vietnam).

Amphidromus zebrinus* (Pfeiffer, 1861)
(Fig. 4b)

Bulimus zebrinus Pfeiffer, 1861: 194.

Amphidromus (Syndromus) zebrinus-Laidlaw & Solem, 1961: 564.

Syndromus zebrinus Schileyko, 2011: 52.

Amphidromus zebrinus Sutcharit et al., 2015: 95, fig. 15-K; Inkhavilay et al., 2019: 95.

Type locality. Siam.

Material examined. VIETNAM, Dong Nai province, 2A, HNUE-LS 000.11, Tan Phu district, Cat Tien Forest, on the way to Bau Sau, 11°27'34.84"N - 107°21'17.43"E, alt. 156 m, 20 May 2021, Nhuong DV leg.

Distribution. Vietnam (Dong Nai, near Binh Dinh, Annam), Laos, Thailand (Schileyko, 2011).

Remarks. The examined specimens share some features with descriptions of *Amphidromus zebrinus* (Pfeiffer, 1861) such as sinistral and conical shell, whorls slightly convex, oblique aperture; simple peristome, moderately expanded.

DISCUSSION

Amphidromus species exhibits remarkable shell morphology diversity. While Thach et al. (2016–2023) have recently published around 140 new taxa from Vietnam, 72 of them have been recognized as synonyms of existing species through combined analysis of morphology and molecular data, such as *Amphidromus thanhhoaensis* Thach & Huber, 2016 = *Aegistohadra roemerri* (Pfeiffer, 1863), and *Amphidromus pankowskiae* Thach, 2020 = *Amphidromus cruentatus* (Morelet, 1875) (Jirapatrasilp et al., 2022; Lee et al., 2022). Given the complexity of these findings, a comprehensive review is essential. Consequently, our study minimally refers to Thach's publications. A comprehensive approach integrating molecular and morphology data is necessary to stabilize the taxonomy of *Amphidromus* species.

Comparatively, Vietnam's *Amphidromus* species count is notable against other nations such as Laos (20 species) (Inkhavilay et al., 2019), Thailand (22 species) (Laidlaw & Solem, 1961; Sutcharit & Panha, 2006), and China (6 species) (Wang & Chen, 2021). Schileyko (2011) reported 23 species in Vietnam. However, our study did not collect specimens of thirteen species due to limitations in survey coverage, notably in Vietnam's central regions. This underscores the necessity for broader sampling to reflect the full spectrum of species diversity.

CONCLUSION

Over the past nearly 15 years, surveys across all three regions of Vietnam have recorded 14 species of *Amphidromus*. In conjunction with Schileyko's (2011) synthesis of 23 species, the current count of *Amphidromus* species in Vietnam has risen to 27, adding 4 species to the Vietnamese fauna: *A. atricallosus perakensis*, *A. leucoxanthus*, *A. perversus natunensis*, and *A. areolabiatus*. This publication provides comprehensive data and illustrations of the 14 collected species. Additionally, it contributes good collections to museums in Vietnam, facilitating easier access to specimens for future research endeavors.

Acknowledgements: This research is supported by the project "Investigate the diversity of aquatic animals as a basis for the conservation of island and cave ecosystems in Ha Long Bay and Bai Tu Long Bay, Quang Ninh province" under grant number UQĐTCB.05/23–25. The fieldwork in Cuc Phuong National Park was funded by the German Federal Ministry of Education and Research (BMBF) under grant number 01DP17052 in the context of the project "Innovative Approaches to Biodiversity Discovery and Characterisation in Vietnam" (VIETBIO); The fieldwork in Phong Nha Ke Bang National park was funded by the Vietnam Ministry of Science and Technology under Grant number ĐTDL.CN-113/21 with the project "Study on biodiversity of Son Doong cave system in Phong Nha - Ke Bang National Park, Quang Binh province for conservation and sustainable development". We sincerely thank Nguyen Van Be, Vo Van Be Hai, Nguyen Thanh Binh, Dinh Phuong Dung, Nguyen Van Hung, Anna Sulikowska-Drozdz for their help in collecting specimens used in this paper. Besides, we would like to express our sincere gratitude to Nguyen Quoc Hung from Phong Nha - Ke Bang NP and the management boards of Phong Nha - Ke Bang NP, Cuc Phuong NP, and Dong Nai Culture and Nature Reserve for their support during our field trips. We also want to acknowledge Mr. Bui Thai Tien for his donation of some specimens and the presentation about his

Amphidromus farm. A big thank you to Parm Viktor von Oheimb, Katharina Viktor von Oheimb, Daniel Mulcahy for comments and suggestions to improve the manuscript.

REFERENCES

- Abbott R. T., 1989. Compendium of landshells. A full colour guide to more than 2,000 of the World's terrestrial shells. Melbourne. American Malacologists, Inc., pp. 240.
- Albers J. C., 1850. Die Heliceen nach natürlicher Verwandtschaft systematisch geordnet. T.C.F. Verlag, Berlin, pp. 262.
- Albers J. C., Martens, E. von., 1860. Die Heliceen nach natürlicher Verwandtschaft systematisch geordnet von Joh. Christ. Albers. Leipzig: Engelmann, pp. 377 [original manuscript of Albers edited and published by Martens].
- Ancey C. F., 1897. On two new species of *Amphidromus*. *The Nautilus*, 11(6): 62–63.
- Barna Pál-Gergely, András Hunyadi, Kurt Auffenberg, 2020. Taxonomic vandalism in Malacology: Comments on Molluscan taxa recently described by N. N. Thach and colleagues (2014–2019). *Folia Malacologica*, 28 (1): 35–76.
- Bavay A., Dautzenberg P., 1909a. Molluscorum terrestrium Tonkinorum diagnoses. *Journal de Conchyliologie*, 56: 229–251.
- Bavay A., Dautzenberg P., 1909b. Description de coquilles nouvelles de L'Indo-chine. *Journal de Conchyliologie*, 57(4): 279–288.
- Collinge W. E., 1903. Report on the non-operculate land Mollusca. *Fasciculi Malayenses Zoology*, 2: 205–218.
- Crosse H., Fischer P., 1863. Note sur la faune malacologique de Cochinchine, comprenant la description des espèces nouvelles ou peu connues. *Journal de Conchyliologie*, 11 (4): 343–379.
- Crosse H., Fischer P., 1864. Faune malacologique de Cochinchine. *Journal de Conchyliologie*, 12: 322–338.
- Eydoux J. F. T., Souleyet L. F. A., 1852. Voyage autour du monde exécuté pendant les années 1836 et 1837 sur la corvette La Bonite commandée par M. Vaillant. Zoologie, Tome Deuxième, Arthus Bertrand, Paris, pp. 664.
- Fischer H., Dautzenberg P., 1904. Catalogue des mollusques terrestres et fluviatiles de l'Indo-Chine 1879–1895. Volume 3. Ed. E Leroux, pp. 390–450.
- Fischer P., 1891. Catalogue et distribution géographique des Mollusques terrestres, fluviatiles & marins d'une partie de l'Indo-Chine (Siam, Laos, Cambodge, Cochinchine, Annam, Tonkin). Autun, pp. 186.
- Fulton H., 1896. A list of the species of *Amphidromus* Albers, with critical notes and descriptions of some hitherto undescribed species and varieties. *Annals and Magazine of Natural History*, 17: 66–94.
- Fulton H., 1901. Descriptions of new species of *Xesta*, *Amphidromus*, and *Cyclostoma* from Madagascar and Perak. *The Journal of Malacology*, 8 (4): 103–104.
- Fulton H. C., 1899. Description of supposed new species of *Streptaxis* and *Amphidromus*. Proceedings of the Malacological Society of London, 3: 302–303.
- Gould A. A., 1843. Description of land mollusks from the province of Tavoy, in British Burmah. *Proceeding of the Boston Society of Natural History*, 1: 137–141.
- Gude G. K., 1903. A classified list of the helicoid land shells of Asia (Part VI). *Journal of Malacology*, 10: 45–62.
- Gude, G. K., 1903. A classified list of the helicoid land shells of Asia (Part V). *Journal of Malacology*, 10: 5–16.
- Haas F., 1934, Neue Landschnecken des Senckenberg museums. *Senckenbergiana*, 16 (2/3): 94–98.
- He J., Zhou Q., 2017. Records of *Amphidromus* from South China. *Shell Discoveries*, 2 (1): 3–7.
- Maitre H., 2007. Rừng người Thượng (Les Jungles mois) (Translate into Vietnamese

- by Luu Dinh Tuan). Knowledge Publishing House, pp. 370. (in Vietnamese).
- Inkhavilay K., Sutcharit C., Panha S., 2017. Taxonomic review of the tree snail genus *Amphidromus* Albers, 1850 (Pulmonata: Camaenidae) in Laos, with the description of two new species. *European Journal of Taxonomy*, 330: 1–40.
- Inkhavilay K., Sutcharit C., Bantaowong U., Chanabun R., Siriwit W., Srisonchai R., Pholyotha A., Jirapatrasilp P., & Panha S., 2019. Annotated checklist of the terrestrial molluscs from Laos (Mollusca, Gastropoda). *ZooKeys*, 834: 1–166.
- Jirapatrasilp P., Huang C. W., Hwang C. C., Sutcharit C., & Lee C. T., 2022. Convergent evolution of *Amphidromus*-like colourful arboreal snails and phylogenetic relationship of East Asian camaenids, with description of a new *Aegistohadra* species (Helicoidei: Camaenidae: Bradybaeninae). *Invertebrate Systematics*, 36 (3): 244–290.
- Laidlaw F. F., Solem A., 1961. The land snail genus *Amphidromus*: A synoptic catalogue. *Fieldiana: Zoology*, 41(4): 507–677.
- Lap V. T., 2009. Natural Geography of Vietnam. Hanoi National University of Education, pp. 345 (In Vietnamese).
- Lee H.G., 2022. Aenigmatic *Aegistohadra* Part 2 (conclusion) -or- An account of a pretty astounding clade of sinistral and enantiomorphic land snail taxa. *Shell-O-Gram*, 63 (4): 3–10.
- Linnaeus C., 1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata [10th revised edition], vol. 1, Laurentius Salvius: Holmiae*, pp. 824.
- Maassen W. J. M., 2001, A preliminary checklist of the non-marine molluscs of West-Malaysia: “a Hand List”. *De Kreukel, Extra Editie 2001*:1–155.
- Mabille J., Mesle G. L., 1866. Observations sur la faune malacologique de la Cochinchine et du Cambodge, comprenant la description des espèces nouvelles (1). *Journal de Conchyliologie*, 14: 117–138.
- Mabille M. J., 1887. Sur quelques mollusques du Tonkin. Bulletins de la Société Malacologique de France. Tome quatrième, 4: 73–164.
- Martens C. E. von, 1864. Diagnosen neuer Heliceen aus dem ostasiatischen Archipel. *Monatsberichte der Königlichen Preussischen Akademien der Wissenschaften zu Berlin*: 523–529.
- Martens C. E. von, 1867. Die Landschnecken. Die Preussische Expedition nach Ost-Asien. *Zoologischer Theil*, 2: 1–447.
- Möllendorff O. F. von, 1894. On a collection of land-shells from the Samui Islands, Gulf of Siam. *Proceeding of the Zoological Society of London*, 1894: 146–156.
- Möllendorff O. F. von, 1898. Die Binnenmollusken Annams. *Nachrichtsblatt der Deutschen Malakologischen Gesellschaft*, 30(5–6): 65–85.
- Möllendorff O. F. von, 1900. Zwei neue *Amphidromus* aus Annam. *Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft*, 32(1–2): 22–24.
- Möllendorff O. F. von, 1901. Zur Binnenmollusken-Fauna von Annam, IV. *Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft*, 33(3–4): 45–50.
- MolluscaBase eds, 2022: MolluscaBase. Accessed at <https://www.mollusca-base.org>
- Morlet L., 1885. Description d'espèces nouvelles de coquilles recueillies par M. Pavie au Cambodge. *Journal de Conchyliologie*, 32(4): 386–403.
- Morlet L., 1889. Catalogue des coquilles recueillies, par M. Pavie dans le Cambodge et le Royaume de Siam, et description d'espèces nouvelles (1). *Journal de Conchyliologie*, 37: 121–199.

- Oesch R. D., Watrous L. E., Barnhart M. C., 2013. Land snails & slugs of Missouri. Missouri Department of Conservation, pp. 214.
- Pfeiffer L., 1857. Descriptions of fifty-eight new species of Helicea from the collection of H. Cuming, Esq. *Proceedings of the zoological Society of London*, 24: 324–336.
- Pfeiffer L., 1857. Diagnosen neuer landschnecken. *Malakozoologische Blätter*, 4: 155–170.
- Pfeiffer L., 1861. Descriptions of new land-shells, in the collection of H. Cuming, Esq. *Proceedings of the Zoological Society of London*, 29: 190–196.
- Pilsbry H. A., 1894. Manual of Conchology, Structural and Systematic, with Illustrations of the Species, Ser. 2, Vol. 9. Academy of Natural Sciences, Philadelphia, pp 1–366.
- Pilsbry H. A., 1900. Manual of Conchology, Ser. 2. Vol. 13. The Academy of Natural Sciences of Philadelphia, Philadelphia, pp. 127–234.
- Reeve L. A., 1848. *Conchologia iconica or, illustrations of the shells of molluscous animals*. Bulimus. London. Pls 37.
- Reeve L. A., 1860. On two new species of shells from Camboja. *The Annals and Magazine of Natural History*, Series 3, 6(33): 203–204.
- Richardson L., 1985. Camaenidae: Catalogue of species. *Tryonia*, 12: 5–49.
- Schileyko A. A., 2011. Checklist of land pulmonate molluscs of Vietnam (Gastropoda: Stylommatophora). *Ruthenica*, 21(1): 1–68.
- Smith E. A., 1893. Descriptions of six new species of land-shells from Annam. *Proceedings of the Malacological Society of London*. 1: 10–13.
- Solem A., 1965. Land snails of the genus *Amphidromus* from Thailand (Mollusca: Pulmonata: Camaenidae). *Proceeding of the United States National Museum*, 117: 615–627.
- Solem A., 1983. First record of *Amphidromus* from Australia, with anatomical notes on several species (Mollusca: Pulmonata: Camaenidae). *Records of the Australian Museum*, 35(4): 153–166.
- Sutcharit C., Ablett J., Tongkerd P., Naggs F., Panha S., 2015. Illustrated type catalogue of *Amphidromus* Albers, 1850 in the Natural History Museum, London, and descriptions of two new species. *ZooKeys*, 492: 49–105.
- Sutcharit C., Naggs F., Ablett J., Sang P. V., Hao L. V., Panha S., 2021. Anatomical note on a tree snail *Amphidromus (Amphidromus) cambojiensis* (Reeve, 1860) from Vietnam (Eupulmonata: Camaenidae). *Journal of Natural History*, 55(17–18): 1059–1069.
- Sutcharit C., Panha S., 2006. Taxonomic review of the tree snail *Amphidromus* Albers, 1850 (Pulmonata: Camaenidae) in Thailand and adjacent areas: Subgenus *Amphidromus*. *Journal of Molluscan Studies*, 72(1): 1–30.
- Sutcharit C., Thach P., Chhuoy S., Ngor P. B., Jeratthitikul E., Siriwit W., Srisonchai R., Ng T. H., Pholyotha A., Jirapatrasilp P., Panha S., 2020. Annotated checklist of the land snail fauna from southern Cambodia (Mollusca, Gastropoda). *Zookeys*, 948: 1–46.
- Sutcharit C., Tongkerd P., Panha S., 2013. First Record on Chiral Dimorphic Population of *Amphidromus inversus annamiticus* (Crosse & Fischer, 1863) from Thailand. *Tropical Natural History*, 13(1): 53–57.
- Tapparone-Canefri C., 1874. *Zoologia del viaggio intorno al globo della regia fregata Magenta durante gli anni 1865–1868. Malacologia (Gasteropodi, Acefali e Brachiopodi)*, Torino, pp. 161.
- Thach N. N., 2016. Vietnamese new mollusks with 59 new species. 48HrBooks Company, Ohio, USA, pp. 205.

- Thach N. N., 2017a. A new species of *Amphidromus* Albers, 1850 (Gastropoda: Camaenidae) from Vietnam. *Gloria Maris*, 55: 123–125.
- Thach N. N., 2017b. Four new species of *Amphidromus* Albers, 1850 (Gastropoda: Camaenidae) from Vietnam. *Gloria Maris*, 56: 35–39.
- Thach N. N., 2017c. New shells of Southeast Asia with 2 new genera and 85 new species. 48HrBooks Company, Ohio, USA, pp. 128.
- Thach N. N., 2017d. Three new species of *Amphidromus* Albers, 1850 (Gastropoda: Camaenidae) from Vietnam, and presentation of *Conus (Rhizoconus) paulkersteni*, nom.nov. pro *Conus kersteni* Thach, 2017 non Tenorio, Afonso & Rolán, 2008. *The Festivus*, 49: 296–300.
- Thach N. N., 2018. New shells of South Asia seashells-freshwater & land snails, 3 new genera, 132 new species & subspecies. 48HrBooks Company, Ohio, USA, pp. 173.
- Thach N. N., 2019. *Amphidromus atricallosus vovanae*, a new subspecies of *Amphidromus* from Vietnam (Mollusca: Gastropoda: Camaenidae). *Gloria Maris*, 57(3): 84–87.
- Thach N. N., 2020. New shells of South Asia. Volume 2. Seashells-Freshwater-Land snails. With one New Genus and 140 New Species & Subspecies, Reply to comments made in error. 48HRBooks Company, Akron, Ohio, USA, pp. 189.
- Thach N. N., 2021. New shells of South Asia and Taiwan, China, Tanzania. Seashells freshwater land snails with 116 new species & subspecies and rejected synonyms, accepted species. 48HrBooks Company, Ohio, USA, pp. 202.
- Thach N. N., 2023. New shells of south Asia and Japan, Taiwan, China with 1 new genus, 1 new subgenus and 123 new species and subspecies including 77 new marine species, 41 new land species 5 new freshwater species and 18 new bivalves. 48HrBooks Company, USA, pp. 141.
- Thach N. N., Huber F., 2014. A new *Amphidromus* (Gastropoda: Camaenidae) from Vietnam. *Basteria*, 78: 35–37.
- Thanh D. N., 2008. Overview on the species composition of the landsnails fauna of Vietnam. *Academia Journal of Biology*, 30(4): 1–15.
- Wang Y. C., Chen Z. Y., 2021. A new species of *Amphidromus* Albers, 1850 (Gastropoda: Stylommatophora: Camaenidae) from China, with notes on *Amphidromus* species. *Folia Malacologica*, 29(1): 33–37.
- Wang Y. C., 2019. Review on the synonymy concerning genus *Amphidromus* Albers, 1850 (Gastropoda: Camaenidae), with descriptions of new species. *The Festivus*, 51(4): 297–305.
- Wang Y. C., 2022. Evaluative notes on the systematics of the “*Amphidromus*” *pervariabilis*-complex and the “*Amphidromus*” *roemerii*-complex (Gastropoda: Stylommatophora: Camaenidae). *Folia Conchyliologica*, 66: 25–32.
- Wu M., 2004. Preliminary phylogenetic study of Bradybaenidae (Gastropoda: Stylommatophora: Helicoidea). *Malacologia*, 46: 79–125.
- Zilch A., 1953. Die Typen und Typoide des Natur-Museums Senckenberg, 10: Mollusca, Pleurodontidae (1). *Archiv für Molluskenkunde*, 82(4/6): 131–140.
- Zilch A., 1960. Gastropoda, Euthyneura. In: Schindewolf O.H. (ed.) *Handbuch der Paläozoologie*, 6: 601–834. Gebrüder Borntraeger, Berlin.