

## Merosesquiterpenes from marine sponge *Smenospongia cerebriformis*

Le Thi Huyen<sup>1</sup>, Dan Thi Thuy Hang<sup>2</sup>, Nguyen Xuan Nhiem<sup>2</sup>, Bui Huu Tai<sup>2</sup>,  
Hoang Le Tuan Anh<sup>2</sup>, Pham Hai Yen<sup>2</sup>, Nguyen Van Dau<sup>1</sup>, Chau Van Minh<sup>2</sup>, Phan Van Kiem<sup>2\*</sup>

<sup>1</sup>Hanoi University of Science, Vietnam National University

<sup>2</sup>Institute of Marine Biochemistry, Vietnam Academy of Science and Technology

Received 31 October 2016; Accepted for publication 11 April 2017

### Abstract

Using various chromatography methods, three merosesquiterpenes belonging to sesquiterpene quinone type, neodactyloquinone (**1**), dactyloquinone D (**2**), and dactyloquinone C (**3**) together with two indole derivatives indole-3-aldehyde (**4**) and indole-3-carboxylic methyl ester (**5**) were isolated from the methanol extract of the Vietnamese marine sponge *Smenospongia cerebriformis*. Their structures were determined by 1D-, 2D-NMR spectra, HR-ESI-MS and in comparison with those reported in the literature.

**Keywords.** *Smenospongia cerebriformis*, merosesquiterpene, sesquiterpene quinone, indole derivative.

### 1. INTRODUCTION

Marine sponges are regarded as a rich source of secondary metabolites with chemically diverse structures and potential biological benefits. Merosesquiterpenes and indole alkaloid derivatives were found to be the main components of sponges, particularly, the genus *Smenospongia*. A huge variety of compounds belonging to these two chemical structure classes have been reported from sponges and possessed a broad range of interest bioactivities, such as antimalarial [1, 2], antimicrobial [1-3], anticancer [3, 4] antidepressant [5], as well as inhibition of the neuronal isozyme of nitric oxide synthase (nNOS) [6]. Herein, we report the isolation and structure determination of three merosesquiterpenes and two indole alkaloid derivatives from sponge *Smenospongia cerebriformis*.

### 2. MATERIALS AND METHODS

#### 2.1. Sponge materials

The sponge *Smenospongia cerebriformis* (Duchassaing & Michelotti, 1864) was collected in Vinh Moc, Quang Tri in August 2015 and identified by Prof. Do Cong Thung, Institute of Marine Environment and Resources, VAST. A voucher specimen (HM08.2015-2) was deposited at the Institute of Marine Biochemistry, Vietnam Academy of Science and Technology.

#### 2.2. General experimental procedures

The <sup>1</sup>H-NMR (500 MHz) and <sup>13</sup>C-NMR (125 MHz) spectra were recorded on a Bruker AM500 FT-NMR spectrometer and TMS was used as an internal standard. Column chromatography was performed using a silica gel (Kieselgel 60, 70–230 mesh and 230–400 mesh, Merck, Whitehouse Station, NJ) or RP-18 resins (30–50 μm, Fuji Silysia Chemical Ltd.), and thin layer chromatography (TLC) using pre-coated silica-gel 60 F<sub>254</sub> (0.25 mm, Merck) and RP-18 F<sub>254S</sub> plates (0.25 mm, Merck).

#### 2.3. Extraction and isolation

Fresh frozen dried samples of sponge *Smenospongia cerebriformis* (15.0 kg) were well grinded and sonicated with hot MeOH three times and then concentrated under reduced pressure to give MeOH extract (SP, 360 g). This extract was suspended in water and then partitioned with CH<sub>2</sub>Cl<sub>2</sub> to give the CH<sub>2</sub>Cl<sub>2</sub> (SPD, 102 g) and water (SPW, 250 g) extracts after removal of the solvents *in vacuo*. Fraction SPD (100 g) was subjected to silica gel column chromatography and eluted with an *n*-hexane - acetone stepwise gradient to give five fractions SPD1 (39.0 g), SPD2 (5.8 g), SPD3 (12.9 g), SPD4 (20.0 g), and SPD5 (2.8 g). SPD2 was chromatographed on a RP-18 column eluting with acetone - water (1.5:1, v/v) to give four smaller fractions SPD2A-D. Fraction SPD2B was subjected

to silica gel column chromatography and eluted with a *n*-hexane-ethyl acetate (1.5:1, v/v) to give compound **4** (ASP2, 11.0 mg). Fraction SPD3 was chromatographed on a silica gel column eluting with *n*-hexane-ethyl acetate (3:1, v/v) to give five smaller fractions, SPD3A-E. Fraction SPD3D (2.2 g) was applied to a silica gel column eluting with *n*-hexane-ethyl acetate (2:1, v/v) to give compounds **1** (ASP16A, 10.0 mg) and **2** (ASP15A, 12.0 mg). Fraction SPD3E (1.8 g) was chromatographed on a

RP-18 column eluting with acetone - water (1:1, v/v) to yield compound **5** (16 mg). Fraction SPD5 was subjected to a silica gel column using dichloromethane-ethyl acetate (10:1, v/v) as eluent to give five smaller fractions, SPD5A-D. Furthermore, fraction SPD5B (0.4 g) was firstly chromatographed on a RP-18 column eluting with acetone-water (2:1, v/v) and then further purified on a silica gel column eluting with dichloromethane - acetone (12:1, v/v) to yield compound **3** (ASP27, 11.0 mg).




Figure 1: Chemical structures of compounds **1-5** from *S. cerebriformis*

**Neodactyloquinone (1):** White amorphous powder;  $\alpha_D^{25}$ : +25.4 ( $c = 0.1$ , in  $\text{CDCl}_3$ );  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ), see table 1.

**Dactyloquinone C (2):** White amorphous powder;  $\alpha_D^{25}$ : +30.2 ( $c = 0.1$ , in  $\text{CDCl}_3$ );  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ), see table 2.

**Dactyloquinone D (3):** White amorphous powder;  $\alpha_D^{25}$ : +21.6 ( $c = 0.1$ , in  $\text{CDCl}_3$ );  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta_{\text{H}}$ : 5.76 (s, H-19), 4.52 and 4.49 (each 1H, br s, H-11), 3.80 (3H, s, 20-OMe), 1.24 (3H, s, H-13), 1.11 (3H, s, H-12), 1.07 (3H, s, H-14);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ , 125 MHz)  $\delta_{\text{C}}$ : 22.0 (C-1), 28.3 (C-2), 32.6 (C-3), 158.6 (C-4), 39.4 (C-5), 31.2 (C-6), 30.4 (C-7), 84.8 (C-8), 37.1 (C-9), 44.9 (C-10), 103.5 (C-11), 20.9 (C-12), 23.1 (C-13), 21.1 (C-14), 27.8 (C-15), 113.9 (C-16), 152.6 (C-17), 181.7 (C-18), 104.9 (C-19), 159.4 (C-20), 181.0 (C-21), and 56.4 (20-OMe).

**Indole-3-aldehyde (4):** White amorphous powder;  $^1\text{H}$ -NMR ( $\text{CD}_3\text{OD}$ , 500 MHz)  $\delta_{\text{H}}$ : 8.04 (s, H-2), 8.13 (d,  $J = 7.5$  Hz, H-4), 7.20 (dd,  $J = 7.5$ , 8.0 Hz, H-5), 7.24 (dd,  $J = 7.5$ , 8.0 Hz, H-6), 7.45 (d,  $J = 7.5$  Hz, H-7), 9.84 (s, H-8);  $^{13}\text{C}$ -NMR ( $\text{CD}_3\text{OD}$ , 125 MHz)  $\delta_{\text{C}}$ : 139.7 (C-2), 120.1 (C-3), 125.7 (C-3a), 122.4 (C-4), 123.6 (C-5), 125.0 (C-6), 113.1 (C-7), 138.9 (C-7a), 187.4 (C-8).

**Indole-3-carboxylic methyl ester (5):** White amorphous powder,  $^1\text{H}$ -NMR ( $\text{CD}_3\text{OD}$ , 500 MHz)  $\delta_{\text{H}}$ : 7.93 (s, H-2), 8.20 (dd,  $J = 3.0$ , 9.0 Hz, H-4), 7.28 (overlapped signals, H-5 and H-6), 7.42 (dd,  $J = 3.0$ , 9.0 Hz, H-7), 3.93 (s, 8-OMe);  $^{13}\text{C}$ -NMR ( $\text{CD}_3\text{OD}$ , 125 Hz)  $\delta_{\text{C}}$ : 133.1 (C-2), 131.0 (C-3), 136.1 (C-3a), 121.6 (C-4), 122.1 (C-5), 123.2 (C-6), 111.5 (C-7), 125.8 (C-7a), 165.6 (C-8), 51.1 (8-OMe).

### 3. RESULTS AND DISCUSSION

Compound **1** was isolated as a white amorphous powder. It had a molecular formula  $\text{C}_{22}\text{H}_{28}\text{O}_4$  which was derived from a pseudo-molecular  $[\text{M}+\text{H}]^+$  ion peak at  $m/z$  357.2039 (calcd. for  $\text{C}_{22}\text{H}_{29}\text{O}_4$ , 357.1988) in the HR-ESI-MS and in conjunction with  $^{13}\text{C}$  NMR data.  $^1\text{H}$  NMR and HSQC spectroscopic analysis of **1** showed the presence of three tertiary methyl groups at  $\delta_{\text{H}}$  0.93, 1.08 and 1.41 (each 3H, s), exocyclic methylene signal at  $\delta_{\text{H}}$  4.54 (2H, br s), methoxy group at  $\delta_{\text{H}}$  3.80 (3H, s) and an olefinic proton at  $\delta_{\text{H}}$  5.74 (1H, s). The  $^{13}\text{C}$  NMR of **1** revealed signals of 22 carbons which were classified by DEPT as nine non-protonated carbons, two methines, seven methylenes, and four methyl carbons.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR disclosed the presence of a dialkoxy-1,4-benzoquinone moiety

Table 1: NMR spectral data for **1-2** and reference compounds

C	$\delta_C^{\#a}$	<b>1</b>		$\delta_C^{\$a}$	<b>2</b>	
		$\delta_C^{a,b}$	$\delta_H^{a,c}$ (mult., <i>J</i> in Hz)		$\delta_C^{a,b}$	$\delta_H^{a,c}$ (mult., <i>J</i> in Hz)
1	21.2	21.2	1.56 (m) 1.77 (m)	78.5	78.5	4.15 (ddd, 5.5, 10.5, 10.5)
2	27.7	27.7	1.27 (m) 1.83 (m)	35.5	35.5	1.80 (m) 2.52 (m)
3	32.7	32.7	2.10 (br d, 14.0) 2.23 (ddd, 5.5, 14.0, 14.0)	30.6	30.7	2.26 (m) 2.40 (ddd, 5.0, 14.0, 14.0)
4	158.3	158.3	-	157.2	157.2	-
5	41.3	41.4	-	40.2	40.3	-
6	30.8	30.9	1.41 (m) 2.00 (ddd, 3.5, 3.5, 14.0)	37.4	37.4	1.64 (m)
7	32.3	32.4	1.58 (m) 1.78 (br d, 14.0)	27.8	27.8	1.53 (m) 1.60 (m)
8	34.6	34.6	-	41.8	41.8	1.39 (m)
9	86.4	86.4	-	37.4	37.4	-
10	47.8	47.8	1.44 (dd, 2.5, 12.0)	60.2	60.2	1.50 (d, 10.5)
11	103.7	103.7	4.54 (br s)	104.1	104.1	4.62 (br s)
12	20.8	20.9	1.08 (s)	21.7	21.8	0.96 (s)
13	24.3	24.3	0.93 (s)	16.4	16.4	1.03 (d, 6.5)
14	19.1	19.1	1.41 (s)	15.0	15.0	0.73 (s)
15	26.8	26.8	2.04 (d, 16.5) 2.72 (d, 16.5)	34.8	34.8	1.99 (d, 14.0) 3.13 (d, 14.0)
16	114.6	114.5	-	130.2	130.3	-
17	151.2	151.3	-	156.5	156.3	-
18	181.5	181.5	-	182.6	183.0	-
19	104.7	104.7	5.74 (s)	105.2	105.2	5.80 (s)
20	159.5	159.5	-	159.0	159.0	-
21	181.5	181.5	-	182.6	182.6	-
20-OMe	56.3	56.4	3.80 (s)	56.4	56.4	3.81 (s)

Measured in <sup>a)</sup>CDCl<sub>3</sub>, <sup>b)</sup>125 MHz, <sup>c)</sup>500 MHz. <sup>#)</sup> $\delta_C$  of neodactyloquinone [8], <sup>§)</sup> $\delta_C$  of dactyloquinone C [9].

( $\delta_H$ : 3.80, 5.74;  $\delta_C$ : 56.4, 104.7, 114.5, 151.3, 159.5, 181.5, 181.5) [7]. The HMBC correlations between H-11 ( $\delta_H$  4.54) and C-3 ( $\delta_C$  32.7)/C-4 ( $\delta_C$  158.3)/C-5 ( $\delta_C$  41.4) suggested an exocyclic olefinic methylene forming at C-11/C-4. Methyl protons H-12 ( $\delta_H$  1.08) have HMBC correlations with C-4/C-5/C-6 ( $\delta_C$  30.9)/C-10 ( $\delta_C$  47.8), indicating the location of a methyl group at C-5. The HMBC correlations between H-14 ( $\delta_H$  1.41) and C-8 ( $\delta_C$  34.6)/C-9 ( $\delta_C$  86.4)/C-10 indicated a methyl group at C-9. The last tertiary methyl group located at C-8 which was indicated by HMBC correlations between proton H-13 ( $\delta_H$  0.93) and carbons C-7 ( $\delta_C$  32.4)/C-8/C-9/C-15 ( $\delta_C$  26.8). The 1,4-benzoquinone moiety linked to sesquiterpene skeleton at C-15 confirmed by HBMC correlations between methylene protons H-15 ( $\delta_H$  2.04, 2.72) and carbons C-7/C-8/C-9/C-13 ( $\delta_C$  24.3)/C-16 ( $\delta_C$  114.5)/C-17 ( $\delta_C$  151.3)/C-21 ( $\delta_C$  181.5). The HMBC correlations from protons H-19

( $\delta_H$  5.74) and methoxy ( $\delta_H$  3.80) to C-20 ( $\delta_C$  159.5) demonstrated for a methoxy group at C-20. Furthermore, carbon chemical shifts of C-9 ( $\delta_C$  86.4) and C-17 ( $\delta_C$  151.3) suggested an ether bridge between C-9 and C-17 which was agreed with molecular formula of **1** C<sub>22</sub>H<sub>28</sub>O<sub>4</sub>. Consequently, structure of **1** was established to be neodactyloquinone, a sesquiterpene quinone previously isolated from the sponge *Dactylosporgia elegans* [8]. Its <sup>1</sup>H and <sup>13</sup>C NMR data were identical with those reported in the literature (table 1) and found to match well [8].

Compound **2** was obtained as a white amorphous powder. The molecular formula of **2** was established as C<sub>22</sub>H<sub>28</sub>O<sub>4</sub> on the basis of HR-ESI-MS (*m/z*: 357.2074, [M+H]<sup>+</sup>; calcd. for C<sub>22</sub>H<sub>29</sub>O<sub>4</sub>, 357.1988) and <sup>13</sup>C-NMR analysis. <sup>1</sup>H NMR and HSQC spectroscopic analysis of **2** showed the presence of two tertiary methyl groups at  $\delta_H$  0.73, 0.96 (each 3H,

s) and a secondary methyl group at  $\delta_{\text{H}}$  1.03 (3H, d,  $J = 6.5$  Hz), exocyclic methylene signal at  $\delta_{\text{H}}$  4.62 (2H, br s), oxygenated methine signal at  $\delta_{\text{H}}$  4.15 (1H, ddd,  $J = 5.5, 10.5, 10.5$  Hz), methoxy group at  $\delta_{\text{H}}$  3.81 (3H, s) and an olefinic proton at  $\delta_{\text{H}}$  5.80 (1H, s). The  $^{13}\text{C}$  NMR of **2** revealed signals of 22 carbons which were classified by DEPT as eight non-protonated carbons, three methines, seven methylenes, and four methyl carbons.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR disclosed the presence of a dialkoxy-1,4-benzoquinone moiety [ $\delta_{\text{H}}$ : 3.81, 5.80;  $\delta_{\text{C}}$ : 56.4, 105.2, 130.3, 156.3, 159.0, 182.6, 183.0] [9]. The HMBC correlations between protons H-11 ( $\delta_{\text{H}}$  4.62) and C-3 ( $\delta_{\text{C}}$  30.7)/C-4 ( $\delta_{\text{C}}$  157.2)/C-5 ( $\delta_{\text{C}}$  40.3) suggested an exocyclic olefinic methylene forming at C-11/C-4. Methyl protons H-12 ( $\delta_{\text{H}}$  0.96) have HMBC correlations with C-4/C-5/C-6 ( $\delta_{\text{C}}$  37.4)/C-10 ( $\delta_{\text{C}}$  60.2), indicating the location of a methyl group at C-5. The HMBC correlations between proton H-14 ( $\delta_{\text{H}}$  0.73) and C-8 ( $\delta_{\text{C}}$  34.6)/C-9 ( $\delta_{\text{C}}$  37.4)/C-10/C-15 ( $\delta_{\text{C}}$  34.8) confirmed the methyl group at C-9. The last secondary methyl group located at C-8 which was indicated by HMBC correlations between H-13 ( $\delta_{\text{H}}$  1.03) and C-7 ( $\delta_{\text{C}}$  27.8)/C-8/C-9. The 1,4-benzoquinone moiety linked to sesquiterpene skeleton at C-15 confirmed by HBMC correlations between methylene protons H-15 ( $\delta_{\text{H}}$  1.99, 3.13) and C-9/C-10/C-14 ( $\delta_{\text{C}}$  15.0)/C-16 ( $\delta_{\text{C}}$  130.3)/C-17 ( $\delta_{\text{C}}$  156.3)/C-21 ( $\delta_{\text{C}}$  182.6). The HMBC correlations from protons H-19 ( $\delta_{\text{H}}$  5.80) and methoxy ( $\delta_{\text{H}}$  3.81) to C-20 ( $\delta_{\text{C}}$  159.0) demonstrated for a methoxy group at C-20. Carbon chemical shifts of C-1 ( $\delta_{\text{C}}$  78.5) and C-17 ( $\delta_{\text{C}}$  156.3) suggested an ether bridge between C-1 and C-17 which was agreed with molecular formula of **2**  $\text{C}_{22}\text{H}_{28}\text{O}_4$ . In addition, the  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR data of **2** were identical with those of dactyloquinone C, a compound also isolated from sponge *Dactylospongia elegans* [9] (table 1) and found to match. Consequently, the structure of **2** was established.

Compound **3** was isolated as a white amorphous powder. The molecular formula  $\text{C}_{22}\text{H}_{28}\text{O}_4$  was deduced on the basis of HR-ESI-MS ( $m/z$ : 357.2039,  $[\text{M}+\text{H}]^+$ ; calcd. for  $\text{C}_{22}\text{H}_{29}\text{O}_4$ , 357.1988) and  $^{13}\text{C}$ -NMR analysis.  $^1\text{H}$  NMR data of **3** also showed the presence of three tertiary methyl groups at  $\delta_{\text{H}}$  1.07, 1.11 and 1.23 (each 3H, s), exocyclic methylene signal at  $\delta_{\text{H}}$  4.49 and 4.52 (each 1H, br s), methoxy group at  $\delta_{\text{H}}$  3.80 (3H, s), and an aromatic proton at  $\delta_{\text{H}}$  5.76 (1H, s). The  $^{13}\text{C}$  NMR and DEPT spectra of **3** revealed signals of 22 carbons including nine non-protonated carbons, two methines, seven methylenes, and four methyl carbons.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR indicated the presence of a dialkoxy-1,4-

benzoquinone moiety [ $\delta_{\text{H}}$ : 3.80, 5.76;  $\delta_{\text{C}}$ : 56.4, 103.9, 114.5, 152.6, 159.4, 181.0, 181.7]. In comparison with **2**, the 1D-NMR spectra of **3** showed the presence signals of an oxygenated tertiary carbon (84.8), a saturated methylene group, and a tertiary methyl group instead of an oxygenated secondary carbon ( $\delta_{\text{C}}$  78.5), a saturated methine group, and a secondary methyl group in the 1D-NMR of **2**, suggesting for the re-arrangement of ether bridge from C-1/C-17 in **2** to C-8/C-17 in **3**. Thus, compound **3** was determined to be dactyloquinone D, a known compound isolated from the sponge *Dactylospongia elegans* [9].




Figure 2: The key HMBC correlations of **1** and **2**

The remaining compounds were elucidated to be indole-3-aldehyde (**4**) and indole-3-carboxylic methyl ester (**5**). Their structures were established based on spectral and chemical evidence, which agreed with previous studies [10, 11].

**Acknowledgement.** This research was supported by Vietnam Academy of Science and Technology under grant number VAST.TĐ.DLB.01/16-18.

## REFERENCES

1. P. Djura, D. B. Stierle, B. Sullivan, D. J. Faulkner, E. V. Arnold, J. Clardy. *Some metabolites of the marine sponges Smenospongia aurea and Smenospongia (.ident.Polyfibrospongia) echina*, Journal of Organic Chemistry, **45**, 1435-1441 (1980).
2. Jin-Feng Hu, John A. Schetz, Michelle Kelly, Jiang-Nan Peng, Kenny K. H. Ang, Horst Flotow, Chung Yan Leong, Siew Bee Ng, Antony D. Buss, Scott P. Wilkins, a. M. T. Hamann. *New anti-infective and human 5-HT<sub>2</sub> receptor binding natural and semisynthetic compounds from the Jamaican sponge Smenospongia aurea*, Journal of Natural Products, **65**, 476-480 (2002).
3. M. L. Kondracki, M. Guyot. *Biologically active quinone and hydroquinone sesquiterpenoids from the sponge Smenospongia sp.*, Tetrahedron, **45**, 1995-2004 (1989).

4. M. -L. Kondracki, A. M. Guyot. *Smenospongine: a cytotoxic and antimicrobial aminoquinone isolated from Smenospongia sp.*, Tetrahedron Letters, **28**, 5815-5818 (1987).
5. Anna J. Kochanowska, Karumanchi V. Rao, Suzanne Childress, Abir El-Alfy, Rae R. Matsumoto, Michelle Kelly, Gina S. Stewart, Kenneth J. Sufka, a. M. T. Hamann. *Secondary metabolites from three Florida sponges with antidepressant activity*, Journal of Natural Products, **71**, 186-189 (2008).
6. E. M. Boyd, J. Sperry. *Synthesis of the selective neuronal nitric oxide synthase (nNOS) inhibitor 5,6-dibromo-2'-demethylaplysinopsin*, Synlett, 826-830 (2011).
7. Hidemichi Mitome, Takahiro Nagasawa, Hiroaki Miyaoka, Yasuji Yamada, a. R. W. M. v. Soest. *A new sesquiterpenoid quinone and other related compounds from the Okinawan marine sponge Dactylospongia elegans*, Journal of Natural Products, **66**, 46-50 (2003).
8. C. Shugeng, G. Zhijie, J. T. Shannon, M. H. Sidney, S. L. John, G. I. K. David. *Marine Sesquiterpenoids that Inhibit the Lyase Activity of DNA Polymerase  $\beta$* , Journal of Natural Products, **67**, 1716-1718 (2004).
9. Hidemichi mitome, Takahiro Nagasawa, Hiroaki Miyaoka, Yasuji Yamada, a. R. W. M. v. Soest. *Dactyloquinones C, D and E novel sesquiterpenoid quinones, from the Okinawan marine sponge, Dactylospongia elegans*, Tetrahedron, **58**, 1693-1696 (2002).
10. M. A. Ashour, E. S. Elkhayat, R. E. R. Ebel, P. Proksch. *Indole alkaloid from the red sea sponge Hyrtios erectus*, Arkivoc, **xv**, 225-231 (2007).
11. Qing-Qing Yang, Marianna Marchini, Wen-Jing Xiao, Paola Ceroni, a. M. Bandini. *Visible-light-induced direct photocatalytic carboxylation of indoles with  $CBr_4/MeOH$* , Chemistry A European Journal, **21**, 18052-18056 (2015).

Corresponding author: **Phan Van Kiem**

Institute of Marine Biochemistry  
 Vietnam Academy of Science and Technology  
 No 18, Hoang Quoc Viet, Cau Giay, Hanoi, Viet Nam  
 E-mail: phankiem@vast.ac.vn; Telephone number: 0983555031.