

The contribution of free radicals in paracetamol degradation by UV/NaClO

Le Truong Giang¹, Dao Thi Phuong¹, Quan Cam Thuy², Dao Hai Yen^{1*}

¹*Institute of Chemistry, Vietnam Academy of Science and Technology*

²*Viet Tri University of Industry (VUI)*

Received 21 November 2017; Accepted for publication 29 December 2017

Abstract

UV/Chlorine is an emerging advanced oxidation process which forms several reactive species including $\cdot\text{OH}$, $\cdot\text{Cl}$, $\cdot\text{OCl}$. This study investigated the contribution of three main free radicals: $\cdot\text{OH}$, $\cdot\text{Cl}$, $\cdot\text{OCl}$ on Paracetamol degradation under different conditions. Benzoic acid (BA), Nitro benzene (NB) and DMOB were used as probe compounds. The second rate constant of $\cdot\text{OH}$, $\cdot\text{Cl}$, $\cdot\text{OCl}$ with PRC were determined: $4.19 (\pm 0.15) \times 10^9 \text{ M}^{-1}\text{s}^{-1}$; $3.71 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$; $3.532 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$, respectively. The formation of free radicals depends on pH. In particular, at pH 5: the contribution of $\cdot\text{OH}$ and ($\cdot\text{OCl}$, $\cdot\text{Cl}$) are 45 %, 41 %, respectively, at pH 8.5, the contribution of free radicals increases up to 63 %.

Keywords. Paracetamol, UV/Chlorine process, reactive species.

1. INTRODUCTION

Advanced oxidation processes (AOPs) are attracting many concerns from researchers in water treatment issues. Their capacity to degrade micro pollutants was demonstrated to be more effective than traditional methods such as: coagulation, filtration, bio-degradation...[1]. NaClO disinfection combined with UV irradiation is effective, low cost consumption and widely applied. UV/NaClO generates primary free radicals $\cdot\text{OH}$ and $\cdot\text{Cl}$ [2] and secondary radicals such as: $\cdot\text{Cl}_2$, $\text{HClO}\cdot$ and $\cdot\text{OCl}$ [3, 4]. $\cdot\text{OH}$ is non-selective oxidation which has ability to react with micro pollutants in the rate constant from 10^8 - $10^{10} \text{ M}^{-1}\text{s}^{-1}$. Chlorine reactive radicals: $\cdot\text{Cl}_2$, $\cdot\text{Cl}$ and $\cdot\text{OCl}$ are selective oxidants. $\cdot\text{Cl}$ prefers to react with substituted aromatics such as: phenol, benzoic acid, toluene and aniline [5]. $\cdot\text{OCl}$ is selective for phenolate and methoxybenzene [5]. The diversity of reactive species in UV/ Cl_2 process contributes to micropollutants degradation in water treatment [3, 4, 6-8]. UV/ Cl_2 process has been applied for many micro pollutants degradation such as: atrazine, carbamazepine, ibuprofen, DEET and caffeine [8, 9], Metoprolol, Triclosan [10, 11]. PRC is a widely used drug all over the world for analgesic and antipyretic agent used for fever, headaches and other minor pain. PRC is one of top 5 most consumed drugs in the world so it presents in

wastewater, surface and even drinking water through human excretion [12, 13]. However, there are no publications about PRC degradation by UV/NaClO.

This study was conducted to identify the kinetic and the relative contribution of 3 main free radicals: $\cdot\text{OH}$, $\cdot\text{Cl}$, $\cdot\text{OCl}$ in PRC degradation by UV/NaClO.

2. MATERIALS AND METHODS

2.1. Chemicals

Paracetamol (PRC) was purchased from Sigma Aldrich and NaOH, HCl, Na_2SO_4 , NaCl, H_2O_2 from Merck, the water distilling machine from Arium Pro, pH measurement machine from Horiba, UV-2900 from Hitachi-Japan. The magnetic mixer and HPLC were obtained from Thermo, Ultra Aqueous C18 ($250 \times 3.2 \text{ nm} \times 5 \mu\text{m}$).

2.2. Experiment

All experiments were conducted in temperature of $25.0 \pm 0.5^\circ\text{C}$ in 2 liters cylindrical reactor equipped with low pressure Hg lamp (UV 254 nm) in the center. UV lamp's photon flow determination was based on H_2O_2 degradation investigations.

The result was $I_0 = 3.41 \times 10^{-6} \text{ Einstein/M}^{-1}\text{s}^{-1}$. PRC concentrations over time were monitored by HPLC.

3. RESULTS AND DISCUSSION

3.1. Determination the rate constant of PRC degradation with $\cdot\text{OH}$

The rate constant of PRC with hydroxyl radical $\cdot\text{OH}$ was determined to evaluate the contribution of $\cdot\text{OH}$ generated in UV/NaClO process. The photolysis process of H_2O_2 was used to generate $\cdot\text{OH}$ following equation:



The effects of pH to PRC degradation were observed and shown in figure 1.




Figure 1: Effects of pH to the rate constant of PRC photolysis reaction

Due to the negligible effects of pH to degradation kinetic of PRC by $\text{H}_2\text{O}_2/\text{UV}$, so all experiments were conducted at unadjusted pH (pH 5). Samples were analysed PRC and H_2O_2 over time.

Based on the experiments conditions conducted, the PRC degradation pathways by hydroxyl radical $\cdot\text{OH}$ follow the pseudo first order kinetic equation and the mechanism was calculated by following equation:

$$-\frac{d[\text{PRC}]_T}{dt} = k_{\cdot\text{OH}/\text{PRC}} [\cdot\text{OH}]_{\text{ss}} [\text{PRC}]_T = k_{\text{obs}} [\text{PRC}]_T$$

$$-\ln \frac{[\text{PRC}]_{T,t}}{[\text{PRC}]_{T,0}} = k_{\text{obs}} t$$

Figure 2 shows that the decrease of PRC over time followed pseudo first order kinetic. Based on the average value of $k_{\text{obs}} = 5.31 \times 10^{-3} \text{ s}^{-1}$ with the assumption that $\cdot\text{OH}$ generated is unvariable, we can calculate the second rate constant of $\cdot\text{OH}$ with PRC = $4.19 (\pm 0.15) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ at pH 5.5-6. This result is higher than in the investigation conducted by Andreozzi to degrade PRC at concentration of (5-20 10^{-6} M); $[\text{H}_2\text{O}_2]_0 < 15 \text{ mM}$ ($k_{\cdot\text{OH}/\text{PRC}} = 2.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ [14, 15] and similar to the result published by Nasma ($k_{\cdot\text{OH}/\text{PRC}} = 4.94 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) [16].




Figure 2: The pseudo first order rate constant of PRC/UV and PRC/UV/ H_2O_2 processes

3.2. Determination the rate constant of PRC degradation with $\cdot\text{Cl}$

To determine the contribution of free radical $\cdot\text{Cl}$, some competitive dynamics experiments were conducted used nitrobenzene (NB) and Benzoic Acid (BA) as probe compounds. The decrease of NB concentration over time mainly caused by $\cdot\text{OH}$ ($k^* \cdot\text{OH} \cdot \text{NB} = 3.9 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$). NB does not react with $\cdot\text{Cl}$, $\cdot\text{OCl}$. Otherwise, BA reacts rapidly with $\cdot\text{OH}$, $\cdot\text{Cl}$ with quite high rate constant ($k^* \cdot\text{OH} \cdot \text{BA} = 5.9 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$; $k^* \cdot\text{Cl} \cdot \text{BA} = 1.8 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$) and does not react with $\cdot\text{OCl}$. All experiments were conducted at pH = 5.5-6 to make sure that the $\cdot\text{OCl}$ concentration is negligible, so we can ignore the contribution of $\cdot\text{OCl}$ in PRC degradation.

Thus, if the assumption of the free radicals $\cdot\text{OH}$ and $\cdot\text{Cl}$ is generated in the system is stable, we can calculate the concentration of $\cdot\text{OH}$ from the competitive kinetic experiment between PRC and NB. The $\cdot\text{Cl}$ concentration was calculated by the concentration reduction of BA in the simultaneous PRC, NB and BA experiments.

For the case of BA:

$$k_{\text{BA}}^{\text{obs}} = k_{\text{UV} \cdot \text{BA}}^{\text{obs}} + k_{\text{NaClO} \cdot \text{BA}}^{\text{obs}} + k^* \cdot\text{Cl} \cdot \text{NB} \times [\cdot\text{Cl}]_{\text{ss}} + k^* \cdot\text{OH} \cdot \text{NB} \times [\cdot\text{OH}]_{\text{s}}$$

Based on the apparent rate constant obtained of PRC, NB and BA in the competitive kinetic reaction, we can calculate the second rate constant of PRC with $\cdot\text{Cl} = 3.71 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$.

3.3. Determination the rate constant of PRC degradation with $\cdot\text{OCl}$

To determine the competitive kinetic, dimethoxybenzene (DMOB) was used as probe compound due to reaction capacity with $\cdot\text{OCl}$ with the second rate constant quite high $2.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$. To facilitate the formation of $\cdot\text{OCl}$ radicals, all

experiments were conducted at pH 8.5, NaClO concentration 100 μM . HCO_3^- 100 mM was added to make sure that all free radicals such as: $\cdot\text{OH}$, $\cdot\text{Cl}$, $\cdot\text{Cl}_2$ are scavenged.

The rate constant of PRC with $\cdot\text{OCl}$ was calculated following this equation:

$$\ln \frac{\text{PRC}_0}{\text{PRC}_t} = \ln \frac{\text{DMOB}_0}{\text{DMOB}_t} \times \frac{k \cdot \text{OCl} \times \text{PRC}}{k \cdot \text{OCl} \times \text{DMOB}}$$




Figure 3: Relationship between degradation rate of PRC and DMOB probe compound

Based on the above results, we can calculate: $k_{\cdot\text{OCl.PRC}} = 1.682 \times 2.1 \times 10^9 = 3.532 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$

3.4. The contribution of free radicals on PRC degradation

Based on the results calculated above, we determined the relative contribution of each oxidizing agent on the PRC total degradation by UV/NaClO. The results were showed on figure 4.

Based on the results showed on figure 4, we can see that the contribution of free radicals on PRC degradation is high. Particularly, at pH 5: the contribution of $\cdot\text{OH}$ and ($\cdot\text{OCl}$, $\cdot\text{Cl}$) are 45 %, 41 %, respectively. However, when pH increases, the concentration of $\cdot\text{OCl}$, $\cdot\text{Cl}$ increases. At pH 8.5, the contribution of free radicals increases up to 63 %.




Figure 4: The relative contribution of free radicals to PRC degradation efficiency by UV/NaClO

4. CONCLUSION

AOPs processes are decided by the rate of free radicals formation. In this study, we determine the second rate constant of 3 main free radicals: $\cdot\text{OH}$, $\cdot\text{Cl}$, $\cdot\text{OCl}$: $4.19 (\pm 0.15) \times 10^9 \text{ M}^{-1}\text{s}^{-1}$; $3.71 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$; $3.532 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$, respectively. This investigation demonstrated that pH highly effects on the formation of free radicals, so their contribution on PRC degradation changes with pH. Particularly, at pH 5: the contribution of $\cdot\text{OH}$ and ($\cdot\text{OCl}$, $\cdot\text{Cl}$) are 45 %, 41 %, respectively and at pH 8.5, the contribution of free radicals increases up to 63 %.

Acknowledgement. This research was supported by National Foundation for Science and Technology Development (NAFOSTED) (grant number 104.06-2013.54).

REFERENCES

- Huerta-Fontela, M., M.T. Galceran, and F. Ventura, *Occurrence and removal of pharmaceuticals and hormones through drinking water treatment*, Water Research, **45(3)**, 1432-1442 (2011).
- Watts, M. J. and K.G. Linden, *Chlorine photolysis and subsequent OH radical production during UV treatment of chlorinated water*, Water Research, **41(13)**, 2871-2878 (2007).
- Fang J., Y. Fu, and C. Shang. *The Roles of Reactive Species in Micropollutant Degradation in the UV/Free Chlorine System*. Environmental Science and Technology, **48(4)**, 334-338 (2014).
- Wu Z., J. Fang, Y. Xiang, C. Shang, X. Li, F. Meng, and X. Yang. *Roles of reactive chlorine species in trimethoprim degradation in the UV/chlorine process: Kinetics and transformation pathways*. Water Research, **104**(Supplement C), 272-282 (2016).
- B. Alfassi, Z., S. Mosseri, and P. Neta, *ChemInform Abstract: Reactivities of Chlorine Atoms and Peroxyl Radicals Formed in the Radiolysis of Dichloromethane*. Cheminform, **20(4)**, 1234-1242 (1989).
- Nam S. -W., Y. Yoon, D. -J. Choi, and K. -D. Zoh. *Degradation characteristics of metoprolol during UV/chlorination reaction and a factorial design optimization*, Journal of Hazardous Materials, **285** (Supplement C), 453-463 (2015).
- Wang, W.-L., Q.-Y. Wu, N. Huang, T. Wang, and H.-Y. Hu, *Synergistic effect between UV and chlorine (UV/chlorine) on the degradation of carbamazepine: Influence factors and Radical Species*. Water Research, **98(7)**, 143-153 (2016).
- Xiang, Y., J. Fang, and C. Shang, *Kinetics and pathways of ibuprofen degradation by the UV/chlorine advanced oxidation process*. Water Research, **90**(Supplement C), 301-308 (2016).

9. Sun P., W. -N. Lee, R. Zhang, and C. -H. Huang. *Degradation of DEET and Caffeine under UV/Chlorine and Simulated Sunlight/Chlorine Conditions*. Environmental Science & Technology, **50(24)**, 13265-13273 (2016).
10. Kong X., J. Jiang, J. Ma, Y. Yang, W. Liu, and Y. Liu. *Degradation of atrazine by UV/chlorine: Efficiency, influencing factors, and products*. Water Research, **90**(Supplement C), 15-23 (2016).
11. Zhou, S., Y. Xia, T. Li, T. Yao, Z. Shi, S. Zhu, and N. Gao. *Degradation of carbamazepine by UV/chlorine advanced oxidation process and formation of disinfection by-products*. Environ Sci Pollut Res, **23(3)**, 1153-1161 (2016).
12. Collado N., S. Rodriguez-Mozaz, M. Gros, A. Rubirola, D. Barceló, J. Comas, I. Rodriguez-Roda, and G. Buttiglieri. *Pharmaceuticals occurrence in a WWTP with significant industrial contribution and its input into the river system*. Environmental Pollution, **185**(Supplement C), 202-212 (2014).
13. Ferrer I., and E. M. Thurman. *Chapter 4 - Analysis of Pharmaceuticals in Drinking Water, Groundwater, Surface Water, and Wastewater*, in *Comprehensive Analytical Chemistry*, M. Petrovic, D. Barcelo, and S. Pérez, Editors, Elsevier, 91-128 (2013).
14. Andreozzi R., V. Caprio, R. Marotta, D. Vogna. *Paracetamol oxidation from aqueous solutions by means of ozonation and H₂O₂/UV system*, Water Research, **37(5)**, 993-1004 (2003).
15. Kosma C. I., D. A. Lambropoulou, and T. A. Albanis. *Investigation of PPCPs in wastewater treatment plants in Greece: Occurrence, removal and environmental risk assessment*. Science of The Total Environment, **466**(Supplement C), 421-438 (2014).
16. Hamdi El Najjar, N., A. Touffet, M. Deborde, R. Journel, and N. Karpel Vel Leitner. *Kinetics of paracetamol oxidation by ozone and hydroxyl radicals, formation of transformation products and toxicity*, Separation and Purification Technology, **136**, 137-143 (2014).

Corresponding author: **Dao Hai Yen**

Biochemistry Department, Institute of Chemistry
Vietnam Academy of Science and Technology
18, Hoang Quoc Viet, Cau Giay, Hanoi, Viet Nam
E-mail: hoasinhmoitruong.vast@gmail.com, dhy182@gmail.com
Telephone: 0985859795.

