

CHEMICAL CONSTITUENTS FROM FRUITS OF *HYDNOCARPUS HAINANENSIS* MERR. (FLACOURTIACEAE) IN VIETNAM

Nguyen Thanh Tra^{1,2}, Truong Bich Ngan¹, Doan Thi Mai Huong¹, Marc Litaudon³,
Nguyen Van Hung¹, Do Thi Thao⁴, Chau Van Minh¹, Pham Van Cuong^{1*}

¹*Institute of Marine Biochemistry - VAST, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam*

²*Institute of Chemistry - VAST, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam*

³*Institute of Natural Product Chemistry, 91190 Gif-sur-Yvette Cedex, France*

⁴*Institute of Biotechnology - VAST, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam*

Received 23 January 2015; Accepted for Publication 18 March 2015

Abstract

Five compounds were isolated from the fruits of *Hydnocarpus hainanensis* Merr. Sleum. (Flacourtiaceae). Their structures were determined by spectroscopic analysis including MS and NMR. The isolates were identified as taraktophyllin (1), hydnocarpic acid (2), 3,4-dihydroxybenzyl alcohol (3), 3,4-dihydroxybenzoic acid (4) and 3-hydroxy-4-methoxybenzoic acid (5).

Keywords. *Hydnocarpus hainanensis*, Flacourtiaceae, cyclopentenoid cyanohydrin glucosides.

1. INTRODUCTION

Flacourtiaceae comprises about 89 genera with 1300 species found throughout the tropical and temperate regions of the world [1]. Genus of *Hydnocarpus* consists of about 40 species, many of them have been used in folk medicine [2]. Previous studies showed that the genus *Hydnocarpus* contains flavonoligans, flavonones, phenolic and acid chaumooric which exhibited antibacterial, antioxidant and anticancer activities [3-7]. In continuation of our research of bioactive compounds from the plants of Flacourtiaceae family, further purification of the crude extract of *Hydnocarpus hainanensis* fruits has led to the isolation of five compounds 1-5.

2. MATERIAL AND METHODS

2.1. General experimental procedures

Optical rotations were recorded on a Polax-2 L polarimeter. Melting points were determined using a Buchi B-545 instrument. ESI-MS were obtained on an Agilent 1100 LC-MSD Trap spectrometer. The NMR spectra were recorded on Bruker 500.13 MHz spectrometer, operating at 500.13 MHz for ¹H

and 125.76 MHz for ¹³C NMR, respectively.

2.2. Plant material

Fruits of *H. hainanensis* Merr. were collected from Quang Tri, Vietnam in November 2006. A voucher specimen (VN-1761) was deposited at the Institute of Ecology and Biological resources, Vietnam Academy Science and Technology.

2.3. Extraction and isolation

Dry powders (0.65 kg) of the fruits of *H. hainanensis* were extracted with ethanol (5 × 1.5 L). The solvents were removed under diminished pressure. The residue (101.7 g) was suspended in water (0.5 L) and then partitioned successively with *n*-hexane, EtOAc. The *n*-hexane, EtOAc and water solutions were concentrated to dryness, affording 43 g, 20 g and 30 g, respectively.

n-Hexane extract (43 g) was fractionated by column chromatography (CC) on silica gel, eluting with *n*-hexane/EtOAc gradient to yield 8 fractions. Fractions 3 (1.7 g) was purified on silica gel CC, eluted with CH₂Cl₂/EtOAc gradient to obtain compound 5 (23 mg).

EtOAc extract (20 g) was subjected to CC on silica gel, eluted with CH₂Cl₂/MeOH gradient to

furnish 7 fractions. Fraction 3 (1.1 g) was purified by CC on silica gel (CH₂Cl₂/acetone gradient) to afford compound **7** (5 mg). Fraction 4 (1.3 g) was separated on silica gel CC (CH₂Cl₂/MeOH gradient), followed by preparative TLC (CH₂Cl₂/MeOH: 95/5) to obtain compound **8** (7 mg). Fraction 5 (1.6 g) was purified on silica gel CC (CH₂Cl₂/MeOH gradient), following by CC on Sephadex LH-20 (MeOH) affording compound **6** (7 mg).

Water extract (30 g) was chromatographed on C-18 (MeOH/H₂O gradient) to give 6 fractions. Fraction 4 (1.1 g) was separated by CC on Sephadex to afford two subfractions. Subfraction 1 (0.6 g) was separated by CC on Sephadex LH-20 following by preparative TLC (CH₂Cl₂/dioxane: 9/1) to give compound **2** (7 mg) and **3** (8 mg). Subfraction 2 (0.3 g) was purified by CC on Sephadex LH-20 (MeOH) to obtain compound **1** (5 mg). Fraction 5 (1.8 g) was separated by Sephadex LH-20 CC (MeOH) yielding two subfractions. Subfraction 2 (0.6 g) was subjected to CC on silica gel (CH₂Cl₂/MeOH gradient), followed by Sephadex LH-20 CC (MeOH) to give compound **4** (5.5 mg). Fraction 3 (350 mg) was separated by CC on silica gel, eluted with *n*-hexane/acetone gradient to give compound **6** (59 mg).

Taraktophyllin (1): Colorless syrup; [α]_D -181 (c 0.083, MeOH). ESI-MS (*m/z*): 310.0 [M+Na]⁺. ¹H-NMR (500 MHz, CD₃OD): 6.16 (1H, dd, *J* = 5.5 and 1.5 Hz, H-2), 6.25 (1H, dd, *J* = 5.5 and 2.0 Hz, H-3), 4.83 (1H, m, H-4), 2.25 (1H, dd, *J* = 4.5 and 14.5 Hz, H-5), 3.06 (1H, dd, *J* = 6.0 and 14.5 Hz, H-5), 4.69 (1H, d, *J* = 8.0 Hz, H-1'), 3.24 (1H, dd, *J* = 7.5 and 9.0 Hz, H-2'), 3.40 (1H, s, H-3'), 3.36 (1H, s, H-4'), 3.36 (1H, s, H-5'), 3.69 (1H, dd, *J* = 5.0 and 11.5 Hz, H-6'a), 3.89 (1H, dd, *J* = 1.5 and 11.5 Hz, H-6'b). ¹³C-NMR (125 MHz, CD₃OD): 82.5 (C-1), 132.9 (C-2), 142.3 (C-3), 74.1 (C-4), 47.9 (C-5), 101.4 (C-1'), 74.7 (C-2'), 77.9 (C-3'), 71.3 (C-4'), 78.2 (C-5'), 62.6 (C-6'), 120.1 (CN).

Hydnocarpic acid (2): Fatty oil [α]_D +36 (c 0.20, CH₂Cl₂). ESI-MS (*m/z*): 253.0 [M+H]⁺. ¹H-NMR (500 MHz, CDCl₃): 5.68 (m, CH-1,2), 2.60 (m, CH-5), 2.34 (t, CH₂-15), 2.30 (m, H-3a), 2.22 (m, H-3b), 2.00 (m, H-4a), 1.38 (m, H-4b), 1.62 (quint, *J* = 7.5 Hz, CH₂-14), 1.26 (8x CH₂). ¹³C-NMR (125 MHz, CDCl₃): 179.2 (C-16), 135.5 (C-1), 130.0 (C-2), 45.6 (C-5), 36.2 (C-6), 33.9 (C-15), 32.0 (C-3), 29.9 (C-4), 29.6-28.0 (C7-C13), 24.2 (C-14).

3,4-Dihydroxybenzyl alcohol (3): Colorless oil. ESI-MS (*m/z*): 140 [M+H]⁺. ¹H-NMR (500 MHz, CD₃OD): 8.02 (d, *J* = 2.5 Hz, H-2), 7.37 (d, *J* = 8.5

Hz, H-6), 7.24 (d, *J* = 2.5, 8.5 Hz, H-5), 4.60 (CH₂-O).

3,4-Dihydroxybenzoic acid (4): White powder. ¹H-NMR (500 MHz, CD₃OD): 7.45 (d, *J* = 1.5 Hz, H-2), 7.43 (dd, *J* = 1.5 Hz and 8.0, H-6), 6.80 (d, *J* = 8.0 Hz, H-5).

3-Hydroxy-4-methoxybenzoic acid (5): Light powder. ESI-MS (*m/z*): 169.1 [M+H]⁺. ¹H-NMR (500 MHz, CD₃OD): 7.58 (1H, d, *J* = 1.7 Hz, H-2), 7.55 (1H, dd, *J* = 8.2 and 1.7 Hz, H-6), 6.84 (dd, *J* = 8.2 Hz, H-5), 3.91 (3H, s, OCH₃).

3. RESULTS AND DISCUSSION

Compound **1** was optically active, [α]_D³⁰ -181 (c 0.083, MeOH). Its ESI-MS (positive) indicated the pseudo-molecular ion peak at *m/z* 310.0 [M+Na]⁺. The ¹H-NMR spectrum of **1** showed the signals of oxymethine protons from 3.24 to 4.69 ppm, two olefinic protons at δ_{H} 6.16 (dd, *J* = 5.5 and 1.5 Hz, H-2) and 6.25 (1H, dd, *J* = 5.5 and 2.0 Hz, H-3), and two protons of a methylene at δ_{H} 2.25 (dd, *J* = 4.5 and 14.5 Hz, H_a-5) and 3.06 (dd, *J* = 6.0 and 14.5 Hz, H_b-5). Analyses of the ¹³C-NMR and DEPT spectra with the aid of the HSQC spectrum of **1** indicated the presence of 12 carbons, including six oxymethines (one anomeric methine at δ_{C} 101.4, C-1'), two methylene groups (one of them was linked to oxygen as indicated by its chemical shifts (δ_{C} 62.6 and δ_{H} 3.69, 3.89, CH₂-6'), an oxygenated quaternary carbon (δ_{C} 82.5, C-1), a double bond and a nitrile functionality (δ_{C} 120.1, C-7). This observation suggested the presence of a sugar moiety. Analysis of the COSY spectrum of **1** revealed two spin-spin coupling systems: correlations of a sugar moiety and a connection starting from H-2 (δ_{H} 6.16) to CH₂-5 (δ_{H} 2.25, 3.06). Analysis of the coupling constants and chemical shifts of the sugar moiety [4.69 (d, *J* = 8.0 Hz, H-1'), 3.24 (dd, *J* = 8.0, 7.5 Hz, H-2'), 3.40 (t, *J* = 7.5 Hz, H-3'), 3.33 (overlapped, H-4' and H-5'), 3.69 (dd, *J* = 5.5, 12.0 Hz, H_a-6'), 3.89 (dd, *J* = 2.0, 12.0 Hz, H_b-6')] determined the presence of glucopyranose moiety in the structure of **1**. In the HMBC spectrum of **1**, the correlations of the protons H-2 (δ_{H} 6.16) and CH₂-5 (δ_{H} 2.25, 3.06) with C-1 (δ_{C} 82.5) and the nitrile carbon C-7 (δ_{C} 120.1) indicated the formation of the cycloheptene ring. The glucopyranosyl moiety was bonded to C-1 as shown by cross-peak of H-1'' (δ_{H} 4.69) with C-1 in the HMBC spectrum. The β -configuration of glucopyranosyl moiety was established by anti coupling constant of H-1'' (*J* = 8.0 Hz). Detailed analyses of the 2D NMR spectra and comparison of the NMR data and optical

rotation with reported values indicated the structure of **1** as taraktophyllin which was previously described [8].

Compound **2** was isolated as colorless oil and optically active, $[\alpha]_D +36$ (c 0.20, CH_2Cl_2). Its ESI-MS indicated the pseudo-molecular ion at m/z 253 $[\text{M}+\text{H}]^+$. In the ^1H NMR spectrum, the signals of two olefinic protons at δ_{H} 5.68 (H-1 and H-2) and the complex overlapped signals of protons in the aliphatic region were observed. The ^{13}C and DEPT of **2** indicated the presence of a carboxylic carbon at δ_{C} 179.2 (C-16), two olefinic carbons at δ_{C} 135.5 (C-1), 130.0 (C-2), a sp^3 methine at δ_{C} 45.6 (C-5), and twelve sp^3 methylenes. Analysis of COSY spectrum of **2** defined the presence of a cyclopentene ring by a connection from H-2 (δ_{H} 5.68) to H-5 (δ_{H} 2.60) via H-2, CH_2 -3 and CH_2 -4. Thus, the remaining signals were assigned to the undecanoic acid side chain. The linkage of C-5/C-6 was determined by correlation of C-6 (δ_{C} 36.2) with H-1 in the HMBC spectrum. Complete analyses of the 2D NMR spectra allowed establishing the structure of **2** as hydnocarpic acid. This compound was previously isolated from several species of *Hydnocarpus* genus [9].

Compound **3** was obtained as colorless oil. The ^1H NMR of **3** displayed the signals of an ABX aromatic system [δ_{H} 8.02 (d, $J = 2.5$ Hz, H-2), 7.37 (d, $J = 8.5$ Hz, H-6), 7.24 (d, $J = 2.5, 8.5$ Hz, H-5)] and two protons of an oxymethylene as singlet at δ_{H} 4.60. These NMR data were in agreement with those of 3,4-dihydroxybenzyl alcohol [10].

The ^1H NMR spectrum of **4** also exhibited the presence of an ABX aromatic system as **3**, forming from three protons at δ_{H} 7.45 (d, $J = 1.5$ Hz, H-2), 7.43 (dd, $J = 1.5$ Hz and 8.0, H-6) and 6.80 (d, $J = 8.0$ Hz, H-5). However, the signal of the oxymethylene was not observed in the ^1H NMR spectrum of **4**. This strongly suggested that the oxymethylene was oxidized into carboxylic acid group. This suggestion was confirmed by comparison of NMR data of **4** with those of previously reported for 3,4-dihydroxybenzoic acid [11, 12].

^1H -NMR spectrum of **5** indicated the signals close to those of **4**, except for the presence of an additional methoxy signal at δ_{H} 3.91. The ABX system was formed by signals of H-2 (δ_{H} 7.58, d, $J = 1.7$ Hz), H-5 (δ_{H} 7.55, dd, $J = 8.2$ and 1.7 Hz) and H-6 (δ_{H} 6.84, dd, $J = 8.2$ Hz). Comparison of the NMR data with reported data in the literature indicated the structure of **5** as 3-hydroxy-4-methoxybenzoic acid [12].

Acknowledgements. The authors thank the Vietnam

Academy of Science and Technology, Vietnam and the National Center for Scientific Research, France (CNRS) for support of the LIA (NAPROCHEMLAB), and Dr. Nguyen Quoc Binh, and MSc Dao Dinh Cuong (VAST- Vietnam for botanical determination.




Figure 1: Structures of compounds **1-5** isolated from the fruits of *Hydnocarpus hainanensis* Merr.

REFERENCES

1. M. W. Chase, S. Zmarzty, M. D. Lledó, K. J. Wurdack, S. M. Swensen, M. F. Fay. *When in doubt, put it in Flacourtiaceae: a molecular phylogenetic analysis based on plastid rbcL DNA sequences*, Kew Bulletin, **57**, 141-181 (2002)
2. M. R. Sahoo, S. P. Dhanabal, A. N. Jadhav, V. Reddy, G. Muguli, U. V. Babu, P. Rangesh. *Hydnocarpus: an ethnopharmacological, phytochemical and pharmacological review*, Journal of Ethnopharmacology, **154**, 17-25 (2014).
3. H. M. Shi, J. Wen, C. Q. Jia, W. Jin, X. F. Zhang, Z. R. Yao, P. F. Tu. *Two new phenolic glucosides from the barks of Hydnocarpus annamensis and their anti-inflammatory and anti-oxidation*, Planta Medica, **72**, 948-950 (2006).
4. S. V. Reddy, A. K. Tiwari, U. S. Kumar, R. J. Rao, J. M. Rao. *Free radical scavenging, enzyme inhibitory constituents from antidiabetic ayurvedic medicinal plant Hydnocarpus wightiana Blume*, Phytotherapy Research, **19**, 277-281 (2005).
5. D. K. Sharma, H. H. Iris. *Hypolipidemic, antiinflammatory and antineoplastic activity and cytotoxicity of flavonolignans isolated from Hydnocarpus wightiana seeds*, J. Nat. Prod., **54**(5), 1298-1302 (1991).

6. J. F. Wang, H. Q. Dai, Y. L. Wei, H. J. Zhu, Y. M. Yan, Y. H. Wang, C. L. Long, H. M. Zhong, L. X. Zhang, Y. X. Cheng, *Antituberculosis agents and an inhibitor of the para-aminobenzoic acid biosynthetic pathway from Hydnocarpus anthelminthica seeds*, *Chemistry & Biodiversity* **7**, 2046-2053 (2010).
7. J. F. Wang, G. F. Yin, X. J. Zhou, J. Su, Y. Li, H. M. Zhong, G. Duan, Y. X. Cheng. *Anti-inflammatory flavonolignans from Hydnocarpus anthelminthica seeds*, *Journal of Asian Natural Products Research*, **13**, 80-83 (2011).
8. D. He, D. Gu, Y. Huang, A. Ayupbek, Y. Yang, H. A. Aisa, Y. Ito. *Separation and purification of phenolic acids and Myricetin from black currant by high speed countercurrent chromatography*, *Journal of Liquid Chromatography & Related Technologies*, **32**, 3077-88 (2009).
9. P. Blaise, M. Fairnes, J. Soulier. *Identification of cyclopentenyl fatty acids by ¹H and ¹³C nuclear magnetic resonance*, *J. Am. Oil Chem. Soc.*, **74**, 727-730 (1997).
10. *Aldrich Library of ¹³C and ¹H FT NMR Spectra*, **2**, 362A; 364C (1992).
11. E. J. Lee, J. S. Kim, H. P. Kim, J. H. Lee, S. S. Kang. *Phenolics constituents from flower buds of Lonicera japonica and their 5-lipoxygenase inhibitory activities*, *Food chemistry*, **120**, 134-139 (2010).
12. S. W. Chang, K. H. Kim, I. K. Lee, S. U. Choi, S. Y. Ryu, K. R. Lee. *Phytochemical constituents of Bistorta manshuriensis*, *Natural Product Sciences*, **15(4)**, 234-240 (2009).

Corresponding author: **Pham Van Cuong**

Institute of Marine Biochemistry, VAST
18 Hoang Quoc Viet, Cau Giay District, Hanoi, Vietnam
E-mail: phamvc@imbc.vast.vn.